

Humanics ErgoSystems, Inc.

Specialists in Ergonomics

ERGONOMICS REVIEW

Balans seating

for VARIÉRUSA

by Rani Lueder, CPE

November 14, 2010

Certificant, Board of Certification in Professional Ergonomics

Humanics ErgoSystems, Inc. | P.O. Box 17388 | Encino, CA 91416-7388 US
Ph: (818) 345-3746 (ERGO) Web: www.humanics-es.com

Ergonomics review: VARIÉRUSA

Table of Contents

Ergonomics review: VARIÉRUSA	1
About this paper	1
Overview (what we have learned).....	2
Movement is important, but it is not enough.....	2
Rethinking “good postures”	3
Rethinking “good postures”	4
Neutral postures	4
Centered postures	5
Sitting versus standing	6
Rethinking postural support.....	6
Forward tilting seats	6
Backrest recline	7
Lumbar supports.....	8
How do workers sit in conventional seats?.....	9
Users rarely take advantage of their chair adjustment features	9
They tend to sit with their back unsupported.....	10
They tend towards forward-leaning positions.....	10
Twisting / rotation / bending movements	11
Balans	13
Balans and lumbar lordosis	13
Balans and EMGs.....	15
Balans and loads on the knees.....	17
Conclusion.....	19
References	21
Additional Research.....	29
About Rani Lueder, CPE	44


Table of Figures

FIGURE 1. The Balans design concept is characterized by an open angle between the thighs and torso utilizing lower leg support to counter the effects of gravity. 1

FIGURE 2. Research indicates that today's adolescents are experiencing high rates of back and neck / shoulder symptoms (e.g., Straker et al, 2008) that often continue into adulthood. Risk is particularly pronounced during growth spurts (Lueder, 2008)..... 11

FIGURE 3. Brunswic (1984a, 1984b) described the added benefit of sitting in the balans chair, related to the integral relationship between the thigh-torso and knee angles..... 15

FIGURE 4. The Thatsit Balans design enables users to promote circulation and reduce leg swelling by activating the venous pumping action of the legs, such as by rocking with one leg on the floor and the other leg on the shin support. 17



About this paper

The following is a general review of the scientific literature relevant to the design and use of the Balans® seat concept - and its comparison with conventional and alternative seating. It was written by Rani Lueder, CPE of Humanics ErgoSystems, Inc. for VARIÉRUSA.

The Balans® seating concept was developed in the 1970's by Hans Christian Mengshoel and designer Peter Opsvik¹ in response to a growing recognition of the limitations of conventional seating from Mandal (1976).

This Balans design is characterized by an open thigh-torso angle and lower leg support that flex (bend) the knees and support balanced movement. Although often referred to as a "kneeling chair" or a "knee-supported chair", the actual locus of lower leg support on the Balans seat is below the knees and at the shin to reduce potential loading at the knees.



Figure 1. The Balans design concept is characterized by an open angle between the thighs and torso utilizing lower leg support to counter the effects of gravity.

Balans chairs are designed with the aim of encouraging neutral postures and facilitating postural changes. Although early versions of the chair provided limited adjustability, subsequent generations incorporated a range of features to include seat depth and height depth adjustability, swivel, back and frontal (chest) support and rocking.

¹ The early versions of the Balans chairs that were based on Mengshoel's design concept were designed by Norwegian designers Svein Gusrud and Oddvin Rykken as well as Peter Opsvik.

Overview (what we have learned)

Long-term sitting in conventional postures increases the risk of developing chronic musculoskeletal disorders, particularly involving the neck, back and shoulders. The rates of spinal disorders in particular have continued to increase in the general population (Harkness et al, 2005) and have become more severe and expensive, affecting our discomfort, health, quality of life and effectiveness at work (Ferguson et al, 2000). Most people will develop severe back injuries and illnesses in their life and once sustained, these tend to recur and worsen.

Movement is important, but it is not enough

Movement matters

We have long known that constrained sitting is bad for our health (s.f., Adams and Hutton, 1983, Duncan and Ferguson, 1974, Eklund, 1967, Graf et. al, 1995, Hunting, et. al, 1980, 1981, Hult, 1954, Langdon, 1965, NIOSH, 1997).

As far back as 1777, Ramazzini described hazards of constrained sitting by writers:

“Now 'tis certain that constant sitting produces Obstructions of the Viscera, especially of the Liver and Spleen, Crudities of the Stomach, a Torper of the Leggs, a languid Motion of the refluent Blood and Cachexies. In a word, Writers are depriv'd of all the Advantages arising from moderate and salutary Exercise.”

Although a range of factors may contribute to back injuries and musculoskeletal disorders, there is clear evidence that long term sitting in awkward and constrained postures greatly increases the associated risk. As today's workers age, they become more susceptible to developing health disorders². At the same time, the work process continues to intensify.

Workers who maintain fixed sitting postures report greater discomfort and chronic disorders (s.f. Graf et al., 1993, 1995). Movement has the potential to reduce these risks (Aaras et al, 2000, Kilbom, 1987).

Yet, as we continue to learn about the causes of musculoskeletal disorders, our focus has also shifted from identifying the best single sitting posture towards a more dynamic view of sitting and movement.

Yet movement is not enough

Postures are also critical. The research clearly establishes that long-term sitting in awkward postures reduce comfort and work effectiveness and introduce health-related risk factor (s.f. review by Pynt and Higgs, 2010). Maintaining conventional slouching postures for only 10 minutes leads to the relaxation of back muscles (Solomonow et al., 2003a; see also review

² Of note, however, the research on back pain and aging is less clear; many young workers experience high rates of back injuries (s.f. review by Dionne et al, 2006)

by Pynt and Higgs, 2010) that transfers associated loads to ligaments and discs (Solomonow et al., 2003c),

Prolonged loading destabilizes the spine (Dolan and Green, 2006; Le et al, 2009; review by Pynt, 2010) and reduces muscle action, increasing laxity of viscoelastic tissues and risk of injury³.

Even when workloads are low, these increase risk of injury following work (Le et al, 2009). Le et al (2009) notes

“Spinal stability refers to the functional and mechanical integrity of its various structures (intervertebral joint, discs, ligaments, facet joint and its capsule, nerve roots, spinal cord, etc.) within their respective physiological ranges.

Poor stability or a deficit in maintaining the various structures within their correct alignment during movement may result in excessive movement and lead to injury (prolapsed disc, facet impingement, nerve root compression, stenosis, etc.) and associated neurological implications of pain, impairment of movement, and substantial loss of work days.”

While attempts to promote movement may mitigate exposure to ergonomic risk factors, it also fostered a misconception that movement is always beneficial and sufficient in itself to prevent harm. The widespread pronouncement that *“the best posture is the next one”* oversimplifies the ergonomics research and is often *flat out wrong* (s.f. review by Lueder, 2005).

All movements are not equivalent; for example, we should strive to avoid postural movements that overload ligaments (Solomonow, 2009), particularly in forward leaning (anterior) and twisted positions (Adams, 1994, 1996a, 1996b; Kumar, 2004; Lueder, 2005) which are commonly evident among today's office workers.

The findings of Solomonow (2009) caused Chaitow (2009) to note:

“Ligaments it seems are far from simply being restraining structures that are strategically placed to support and stabilize joints... They are sensory organs that provide proprioceptive input to the CNS, as well as having reflexive influences on associated muscles, which therefore become major elements in the stabilization of joints”.

In conclusion, movement is critical for our well-being, but it is not the only consideration; the postures that we assume as we move are also paramount. Taken to extremes, a strict emphasis on promoting movement for its own sake may even introduce new risk factors.

³ Le et al (2009)'s findings involved static work of up to 3 hours is associated with an increased laxity.

Rethinking “good postures”


Neutral postures

We are also rethinking the concept of *neutral postures* (Ankrum and Nemeth, 2000; Claus et al, 2009; Reinecke et al, 1994b). Of course, researchers recognize the benefits of *neutral postures*, which equalize the loads over the body and reduce the physical demands associated with range of motions. However, we continue to debate what *neutral postures* represent and how these might vary between individuals and changes in position.

Some research approaches, particularly muscle activity (EMGs) often lead to confounded conclusions (Smoliga et al, 2010; Ankrum 2000a; Ankrum 2000b). Much of what we know from the existing research is based on flawed or limited research approaches that leave some key findings in question.

For example, most findings are based on short-term studies, despite overwhelming evidence to the consequences of constrained sitting⁴ (Beach et al, 2005). Research often measured a limited set of variables in unrealistic short-term laboratory environments.

Measurements of contours of the spine are often been influenced by external tissues of the users. Further, research has failed to adequately consider the implications of the dramatic increase in obesity in the American population⁵; research findings are largely based on people who are young and fit and free of musculoskeletal disorders, age-related degeneration of the spine (including stenosis), nervous system (Bazzucchi et al, 2005; Mariconda et al, 2007) and circulatory disorders that cause edema.



Further, the implications individual differences are poorly understood. For example, females have been found to have greater lumbar lordosis among adolescents (Straker et al, 2008) and adults (Dunk and Callaghan⁶, 2005; de Carvalho et al, 2007).

⁴ Particularly on viscoelastic creep of ligaments and tendons and discs.

⁵ The Centers for Disease Control estimates that in 2006 that 67% of the non-institutionalized US adult population is overweight or obese; 34% are obese. www.cdc.gov/nchs/fastats/overwt.htm

⁶ Dunk and Callaghan (2005) found that females had greater lumbar lordosis and a slightly forward leaning spine compared to males when sitting. Females also positioned themselves closer to the front of a seat pan.

Centered postures

One important benefit of neutral postures is that it enables users to maintain position while sitting close to their center of gravity. Indeed, postures that shift away from these neutral positions are tolerated poorly and for less time (Reinecke et al, 1994b)

It is often difficult to sit upright and unsupported for very long in conventional seating. Most people would rather slump than perform the muscle work needed to sit upright. Leaning against a backrest reduces both intradiscal pressures in the spine and loads at the back portion of the spine (fixator loads involving the facet joints) relative to conventional upright postures (Rohlmann et al, 2001)

Several things happen when we maintain conventional unsupported postures – even with movement. Of particular importance, we tend to slump forward, reversing the lumbar curve (lumbar kyphosis) (Bridger et al, 1989)

If users are fit, their strong abdominal muscles might help stabilize their postures (Corlett and Eklund, 1984) although this is not fully agreed on (Kumar, 2004). Fit or not, postural support shifts from the muscles to the ligaments that support the spine. Ligaments deform, increasing risk of spine and joint injury.

Sitting in conventional chairs generally shifts the user forward from their center of gravity. Users benefit when the center of rotation of the chair is close to that of the user. Centered positions facilitate changes of posture (Andersson, 1986, Fleischer et al, 1987). Andersson (1986) concluded that the relationship between the pivot point of the chair and its user is more important than the dimensions of the backrest^{7, 8}.

Conventional chairs that recline by tilting back at the knees are particularly problematic in this respect. During recline, standard knee-pivoting chairs take the user back, down and away from their worksurface. Chairs that recline in this way will correspondingly expose users to additional risk as they become obliged to elevate their arms and extend their reach and their neck to see their visual target (typically, the document or computer screen).


⁷ Corlett (2002) and Rebiffe (1980) suggested the optimum height of the lumbar support depends on the users' activities. For example, the driver of a car benefits from a lumbar support that is higher than that of the passenger, as driver's arms are higher (on steering wheel), and must reach the controls.

⁸ A close fit between users and their chairs' center of rotations also helps prevent the "shirrtail effect", where the backrest displaces upward, pulling up users' shirrtails.

Sitting versus standing

Until recent years, it was assumed that standing is better for our spine than sitting.

Upon sitting, our hamstrings flex, rotating the pelvis back and excessively flattening the lumbar curve. On the other hand, standing causes the pelvis to rotate forward, thereby increasing the extent of lumbar lordosis.



Research that is more recent suggests that both standing and sitting are problematic. Standing postures are associated with greater pelvic tilt and lumbar lordosis (De Carvalho et al, 2010, Keegan, 1953).

Early findings (Andersson et al 1974, 1975, 1986) that loads on the spine are greater when sitting than when standing have since been contradicted by studies using more sophisticated technology (Wilke, 1999, Solomonow and others), suggesting that some of these early conclusions were limited by the technology at the time. However, Leivseth and Drerup (1997) found less spinal shrinkage when sitting (particularly relaxed sitting) than with standing work. They attributed this to several factors, but particularly to the greater bending and twisting while standing.

Rethinking postural support

Forward tilting seats

Much of this shift in focus regarding the adverse health effects of long-term constrained sitting resulted from the writings of Mandal (1976, 1981, 1982), a Danish plastic surgeon. He associated conventional sitting with a flattening of the lumbar spine.

While Mandal generated considerable interest in the notion of promoting a more natural and healthful sitting posture by tilting the seat pan forward, many noted that users found it difficult to maintain this posture without supporting the lower legs because of the greater effort required to prevent sliding forward / resisting the forward gravitational slide. For this reason, the conventional forward tilting seat is generally considered a failed concept – particularly with the substantial forward tilts Mandal recommended 15 or 20 degrees. Those have been shown not to work because it takes so much more muscle work to sit in that position⁹.

⁹ Even so, this writer has sometimes found forward sloping seat designs that are fairly effective. The basis for these differences are not obvious but seem to relate to the interaction between the center of gravity of the seat versus that of the user, and the ability to establish an open posture that does not

Forward tilting seats that lack leg support did not significantly affect the negative impact of forward reaching (Bendix et al 1988b¹⁰).

Bridger (1988) provides an excellent explanation for why forward-sloping seats might prove less effective than anticipated.

Bendix and Biering-Sorensen (1983) evaluated subjects' postural adaptations to a conventional seat that could be adjusted to slope forward at angles of 0, 5, 10 and 15°. They found that with increasing forward slope, the spine moved forward toward lumbar lordosis.

However, in comparing the 0 and 15° seats, an increase in lumbar curvature of only 4° was observed. This occurred because subjects adapted their sitting posture to the sloping seats by extending the hip joints. The increased trunk-thigh angle brought about by the forward-sloping seat did not lead to a straightforward decrease in lumbar flexion...

As these authors point out, postural adaptation to a forward sloping seat may take place in a number of ways. The whole body may tilt forward, in which case no increase in the trunk-thigh angle need occur. Alternatively, the hip joints may extend, thereby increasing the trunk-thigh angle but without necessarily altering the posture of the spine and pelvis. Finally, as suggested by Mandal, the increased trunk-thigh angle may be accompanied by anteriorpelvic rotation, which would result in a reduction in lumbar flexion, given that the upper trunk did not move. These possibilities represent extremes and might arguably occur in combination.

Backrest recline

Leaning back provides important benefits.

Backrests perform several important functions. These supports may

- a) open the thigh-torso angle;
- b) transfer physical loads from gravity onto the backrest;
- c) sometimes promote lumbar lordosis (Andersson et al, 1986); and
- d) help stabilize the spine (Bendix et al, 1996),

require them to exert a great amount of effort . Often, those seats that work stabilize postural support (such as with the Balans) or provide some sort of a rocking motion.

¹⁰ It should be noted that this short term study examined forward tilting seats but NOT Balans-type chairs. Of note, the study found that users still preferred higher seats that tilt forward.

Backrests may both reduce loads on the spine (intradiscal pressure) and muscle work (Andersson et al, 1974, Yamaguchi and Umezawa, 1970 with reclined seat pans).

These supports stabilize posture by reducing the effort required to fight gravity.


- a) as the weight of the torso shifts back against the backrest, and
- b) as the angle between torso and the legs increases.

Yet reclining also has disadvantages.

Many (perhaps even most) intensive computer users slump against their backrest, locking in their pelvis and causing them to lose (or reverse) their lumbar curve (Dolan and Adams, 2001).

Corlett and Eklund (1984) note, “this will lead to increased pressure on and within the discs, both from forces arising from the stretched muscles and ligaments and the increased wedging at the anterior [forward] edges of the disks”. Bendix (1996) suggested, “The traditional conception that a backrest facilitates lordosis is apparently not true”.

There are also functional limitations associated with reclining. Loads on the shoulders and arms may increase when reclining causes the users to move back against their work items. It is difficult to lean back when our visual target is a document and our hands need to reach the mouse. Reclined postures increase loads on the neck as employees attempt to meet the visual field requirements of the task (Grandjean et al, 1983, Corlett, 1999).



One example of a reclined posture with backrest support (Corlett and Eklund, 1984, with permission)

Lumbar supports

Research indicates that lumbar supports can reduce load on the spine (Andersson et al, 1974b, 1975). By tilting individual vertebra, it also increases pressures at the front of the discs (Adams et al, 1996a, Bendix et al, 1996, Corlett, 1999).

Although Andersson's research findings suggest that lumbar supports can reduce intradiscal loads on the lumbar spine, the benefits of backrest lumbar supports are not consistent (Corlett, 1999). Bendix et al (1996) reported that lumbar supports on backrests helped to reinstate lumbar curves compared to straight backrests while performing tasks, but not during passive sitting and reading. Using pig cadavers, Brodeur and Reynolds (1990) concluded that lumbar supports have little effect on the contours of the lumbar spine. Rather, they found that the lumbar curvature is primarily affected by the pelvic angle.

Characteristics of the lumbar support vary between users¹¹ and may vary over time for the same individual. Corlett (2002), Rebiffe (1980) and others suggested that the optimum height of the lumbar support depends on the users' activities. For example, the driver of a car would benefit from a higher lumbar than the passenger where the driver's arms are higher (on the steering wheel), and they need to reach the controls.

Additionally, lumbar supports only benefit users if they are properly designed, correctly adjusted for the user, and the user sits in the chair in a manner that takes advantage of the feature.

How do workers sit in conventional seats?

Users rarely take advantage of their chair adjustment features

In her review of the research, Lueder (1994, 2005) summarized a broad range of studies indicating that most users do not adjust their chair or take advantage of key seat features. Vink et al (2007) reported that many or most workers never adjust their chair; adjustments performed are frequently limited to the seat height.

¹¹ Pregnant women and heavy users, for example, have more forward center of gravities. Tichauer (1978) notes that while men have centers of gravity above their hip socket, for women these are forward. Perhaps this is why various researchers (s.f. Bridger et al, 1989) have reported that women have deeper lumbar contours than men.


They tend to sit with their back unsupported

People generally prefer sitting to standing.

Unfortunately, when doing so for extended durations in constrained positions, their postures become increasingly hazardous.

In today's offices many or most of office employees work with their backs unsupported by their backrest (e.g., Dowell et al, 2001). In his presentation, Dowell (2001) reported that the large majority of workers work with their back unsupported at any given time, although the rates vary with the task.

Please take a walk around your office to verify this for yourself. This writer once surveyed a large telecom center with over 1,000 intensive computer users. There was not a single person leaning back against their backrest in the entire center.



They tend towards forward-leaning positions

Unsupported sitting posture (Corlett & Eklund, 1984, with permission)

Sitting in static positions leads to small movements in our chair that we often refer to as “fidgeting” (Bhatnager et al, 1985, Fenety et al, 2000). That is, as people become uncomfortable they tend to move more, perhaps to compensate for reduced effectiveness at work. This tendency to fidget with time is associated with a tendency towards forward oriented (anterior) movements and postures that increase loads on the spine and soft tissues¹² (Andersson, 1980, Andersson et al, 1974a, 1975, 1986, Bhatnager et al, 1985, Rohlmann et al, 2001, Wilke et al, 1999).

In fact, loads on the spine (intradiscal pressures) are almost twice as high when flexing forward during unsupported sitting – and almost three times higher than relaxed sitting (Wilke et al, 1999). Others have found that leaning forwards (flexion) led to a dramatic increase in intradiscal pressures as well (Andersson et al, 1974a, 1975, Rohlmann et al., 2001).

¹² Rohlmann et al (2001) suggests that different conclusions might be attributed to the complex geometry of the spine and how much we do not know. For example, high intradiscal pressures in the spine reflect greater loads on the forward portion of the spinal column, but tell us little about loads transferred by the bony protrusions (facet joints) of the spine.

Forward-leaning postures also increase risk of disc rupture because the posterior segment of the disc lacks the strength to bear the associated forces – even when intradiscal pressures are low¹³ (Adams et al, 1994).

The posterior longitudinal ligaments are considerably thinner than the anterior (front) ligaments. Further, the fibrous tissues that surrounds the intervertebral discs¹⁴ are not equivalent – anterior postures are much likely to cause tearing.

Fleischer (1987) emphasized the importance of designing chairs to enable users to promote movements that are known to be beneficial, particularly in the fore-aft / upright-reclined directions (as opposed to forward-leaning / anterior postures).

Twisting / rotation / bending movements

Twisting is common; in fact, it is difficult to function without it. Twisting is also unique to human beings (Kumar, 2004). Twisting (i.e., axial rotation of the spine) increases compressive loads on the spine (Au et al, 2001) and risk of injury (Au et al, 2001, Kumar et al., 1998, Kumar, 2001).



Figure 2. Research indicates that today's adolescents are experiencing high rates of back and neck / shoulder symptoms (e.g., Straker et al, 2008) that often continue into adulthood. Risk is particularly pronounced during growth spurts (Lueder & Rice, 2008).

A key concern is how we can teach these young people to avoid risk and learn proper sitting habits that they can take with them into adulthood.

Kumar (2001) described spinal rotation as “a destabilizing motion for an inherently unstable structure”. He explained how prolonged and extreme twisting could damage joints. First, heightened loads increase the forces acting on the joints, deform connective tissues and ultimately destabilize the joints. With time, as muscles fatigue and joints weaken, the resulting imbalance can lead to unnatural and uncoordinated movements at the joints that can result in injury.

Research suggests that extremely small (less than 2° per vertebral segment) rotations are not harmful

and may even benefit users¹⁵ (Van Deursen et al, 2001). Such micro-movements correspond to the natural / free range of motion of the individual motion segments that make

¹³ These researchers noted that serious disc failure is closely associated with the “moment arm” forces associated with forward bending – even when the compressive loads on the spine itself are not particularly high. They add, “Conversely, if the bending moment is small or absent, no amount of compression can damage the soft tissues before the vertebrae”.

¹⁴ This fibro-cartilaginous tissue of the intervertebral discs is called the annulus fibrosus.

up the spine¹⁶. Minute rotary movements¹⁷ in one's chair may reduce forces acting on the lumbar spine from improved disc' nutrition and lessened back pain (Lengsfeld et al, 2000b).

Yet our spine enables us to twist considerably farther than the 2° mentioned above – actually (when including motion of each of the segments), up to about 70° to each side across the spine (Kumar, 1996, 2004).

Kumar (2004) described the massive body of research demonstrating a very strong relationship between twisting and back injury¹⁸. Some suggest that as little as 20° of twist involving across the mid-back may greatly increase the risk of disc herniation. Kumar described a number of possible mechanisms for this increased risk of injuries, including compression of spine.

Yet biomechanical forces on the spine (moment arm) alone cannot explain the increase in risk with twisting. Au et al (2001) found that even when biomechanical loads are equivalent in different postures, twisting resulted in considerably greater compression of the spine than leaning forwards or controlled sideways bending. The authors note, "it is interesting to consider that the torso is very limited in the production of dynamic twisting torque", even when relatively small levels of force are involved.

Kumar (2004) describes the effect of twisting as jamming the facet joints (bony protrusions of the vertebra), twisting intervertebral discs, tightening some ligaments while slackening others.

Even so, the associated risks vary across the spine and are greater at the lumbar spine. The different vertebrae that make up the spine are very flexible, and designed to rotate to different degrees. Part of this greater flexibility of the lumbar spine is related to the facet joints, which are differently designed to prevent rotation and lateral bending (flexion) displacement (Evjenth and Hamberg, 1985)

Seated twisting also reduces muscle strength. In his previous research, Kumar (2004) found that sitting forward in a neutral posture requires the least amount of strength, but the loads increase as the user moves to 20° of combined vertebral rotation. He concluded, "Thus, when it comes to forceful exertions involving axial rotations, human capability is considerably limited". He continues that "with increasing reach distances, the strengths significantly declined..."

Such studies point to twisting – even rotation while seated at the low (lumbar) and mid-back (thoracolumbar junction) may sometimes increase risk substantially.

¹⁵ This laboratory study by Van Deursen et al (2001) used pig cadavers. These findings may well have been confounded because they removed the facet joints (bony extensions at the back of vertebra) and related spinal components. Even so, these researchers maintained that this did not affect results as the range of rotation was within the joint's free inter-space range of movement.

¹⁶ Panjabi and White (2001) suggest the natural range may be closer to 3°.

¹⁷ These researchers used one patient diagnosed with degenerative instability of the lumbar spine. The rotational movements of the chair were 1.2° to the right and left, at a frequency of .22 Hz.

¹⁸ Risk increased even when lifting was not involved. For example, Kumar (2004) describes research by Marras (1993) that found "twisting without lifting is associated with disc prolapse with an odds ratio of 3.0. A combination of twisting and lifting increased the odds ratio to 6.1.

We need more research to understand the full impact of these issues. Even so, caution is warranted, we expect people to rotate sideways to some extent during seated work, but then it should not be encouraged either.

Balans

Balans and lumbar lordosis

Although the research findings varied, the majority demonstrated improvements in lumbar lordosis – that is, a shift towards more neutral postures - when sitting on the Balans chair.

Further, the flexed knee position may work in concert with the increased thigh-torso angle in promoting neutral postures. The Balans chairs tested in these studies lacked the backrests currently available in some models of the Balans chair.

Adams et al (1990)¹⁹ compared the contours of the spine associated when sitting on the Balans and conventional chairs as well as standing and lying. These researchers reported that sitting on the Balans chair was associated with the most “neutral”. They suggested that these benefits would be particularly beneficial for some users, given age-related changes in muscle, body fat composition, bone density, loss of elasticity of connective tissues.

The gradual age-related dehydration of vertebral discs will reduce these users’ ability to withstand compression and misalignment of the spine.

Frey and Tecklin (1986) found that the lumbar lordosis of users sitting in a Balans Multi-Chair was more neutral than users sitting in a conventional chair, and bore a closer resemblance to the relaxed posture while standing at a workstation.

Bendix et al²⁰ (1988a) found that subjects adapted to a Balans chair exhibited more forward tilt of the pelvis and lumbar lordosis than did users sitting on a chair with a forward-tilting seat pan. Balans users also sustained a more upright head and trunk posture. Although these researchers noted that users generally tended to prefer the forward tilting chair over longer periods, one wonders whether this preference might have related to a greater ease of slumping in the forward tilted chair. The authors suggested that the Balans chair might be a good alternative for some situations.

¹⁹ Adams et al (1990) used a 3D digitizer in this pilot study to evaluate the cervical, thoracic, lumbar, and sacral angles relative to a vertically projected line. For one experimental group, these researchers compared the spinal profile on the Balans and other conventional seats with limited adjustments; for the other experimental group, they compared the profiles associated with standing and lying supine (face up) and prone (face down).


²⁰ These researchers used a stadiometer to evaluate spinal shrinkage of 12 healthy subjects over 3 weeks while performing office and simulated assembly work. Spinal shrinkage was measured over each sitting period; this measure is considered an index of compressive loads acting on the spine.

Bennett et al (1989) measured muscle activity/ EMGs and lumbar lordosis²¹ of 20 subjects sitting in static and dynamic postures in Balans Multi, straight-backed and conventional office chairs and while standing. These researchers found that sitting in the Balans chair increased lumbar lordosis relative to sitting in upright and conventional seating, though less so than when standing.

Bridger et al (1989) measured the spinal angles and hip and lumbar mobility of 25 female subjects standing and in four sitting postures. Lumbar lordosis decreased with reductions in the trunk-thigh angles, which also related to a reduction in hip mobility.

Ericson and Goldie (1989) compared the extent of spinal shrinkage associated with sitting on a conventional, forward-sloping²² and Balans chairs. The subjects' spines shrank more when sitting on the Balans than on conventional chairs. The authors attributed this difference to the lack of a backrest, emphasizing the importance of providing back support with the Balans.

Bishu et al (1991) reported inconsistent results with their small pilot study involving short-term use of folding metal chair, the classroom chair, and the Balans. Findings from the small size and short-term 30 minute sitting trials were inconclusive. Although the Balans was not found most comfortable, the researchers conclude that chair designers need to focus more on the included angle between the seat pan and backrest as well as on the shape of the backrest.



²¹ These researchers used a flexicurve "flexible ruler" to measure lumbar lordosis. Subjects aged between 22 and 37.

²² The forward sloping chair was the "Ullman chair" with front half of seat sloping forward and back half horizontal. Eight healthy subjects worked on computers at their workplaces in 45 minute trials. Eight healthy subjects used to professional VDU work volunteered to participate in this field study, which was performed at their own workplaces.

Bettany-Saltikov et al (2008) compared the lumbar curvatures of users²³ sitting on "kneeling" with conventional computer chairs. They concluded the use of "kneeling chairs" set with 20° of forward tilt did indeed promote neutral lumbar curvatures that more closely resembled standing postures.

Brunswic (1984a, 1984b) found that users sitting on the Balans seat evidenced improved lumbar lordosis in the experimental setting, but the benefits disappeared in the real world setting. This writer presumes the loss of benefits related to the non-

adjustable working height of the users. She emphasized the importance of ensuring the seating accommodated the task and the work environment.

Link et al (1990) compared the lumbar curves of subjects standing and in a conventional and Balans Multi-Chair²⁴. Subjects had significantly more lumbar lordosis / extension when sitting in the Balans Multi-Chair compared with sitting in the conventional chair.

Balans and EMGs

A number of EMG studies have found that muscle loads increased when sitting on the Balans chair. None of these studies compared sitting on Balans chairs that provide back support with conventional seating. Further, these studies have been closely controlled and lack the real world settings that today's office workers operate in.

Yet EMG research is controversial and often lead to confounded conclusions (Smoliga et al, 2010; Ankrum 2000a; Ankrum 2000b). The goal is not to minimize EMGs – after all, exercise is beneficial to health - but rather to avoid excessive muscle loads that exceeds the users' ability to recover. Further, Ankrum points out that much of the research has confused statistical significance with meaningful differences.

Bennett et al (1989) measured muscle activity/ EMGs noted that, given that lumbar lordosis improved in the Balans chair, the findings that Balans sitting involved higher EMGs than sitting in conventional chairs "suggests that EMG activity is not a good indicator of changes in lumbar posture". That is, changes in EMG activity are unable to evaluate the benefits associated with improvements in lumbar lordosis.

Figure 3. Brunswic (1984a, 1984b) described the added benefit of sitting in the Balans chair, related to the integral relationship between the thigh-torso and knee angles.

She noted "*The effect of moving the hips and knees was additive in a 1:2 proportion such that an increase extension of the knees of 20° correspond roughly to a 10° flexion of the hips.*"

[That is, when sitting in the Balans, bending the knees promotes lumbar lordosis, and the benefit represents half that of the increased thigh-torso angles.]

The simultaneous alteration of these two components could flex the spine in proportion to the addition of each phenomenon taken separately."

²³ These researchers used a portable 3D mechanical digitizer "MIDAS System" to assess lumbar curvature, while placing anatomical landmarks on the 20 subjects, aged 18-35.

²⁴ Sixty-one men between 20 and 30 years of age served as subjects. Lumbar curves were measured with a flexible ruler with subjects standing and then sitting in the two chairs.

Shenoy and Aruin (2007) compared the EMGs associated with sitting on a forward-tilted seat and the Balans Multi Chair without back support²⁵. Sitting on both the Balans and standard forward-tilting seat pan resulted in anticipatory activation of trunk and upper leg muscles in some postures. These findings suggest that although the forward-tilting seat and semi-kneeling body position might help in preserving a normal lordosis, it is not associated with anticipatory activation of lower leg muscles, which might reduce the ability of an individual to counteract muscle forces associated with these postures.

Lander et al (1987) found that EMG levels were higher after sitting on a Balans chair without back support, which presumably related to the work configuration and lack of backrest²⁶. Cram and Vinitzky (1995) emphasized the importance of providing pelvic and back support with alternative seating.

²⁵ . Nine healthy subjects were seated on either the Balans or a conventional chair with arms extended. They were instructed to initiate body exertions in four directions.

²⁶ Twenty healthy subjects were randomly assigned to one of two groups. Group 1 subjects sat in the Balans chair for a 30-minute study period and then sat in a conventional office chair for an additional 30-minute period. Group 2 subjects were studied in the reverse seating order.

Balans and loads on the knees

Some studies point to a lack of any evidence of negative loading on the knees when sitting on the Balans. Two very early studies suggested that some users reported discomfort at the knees when sitting on Balans chairs. Both of these were poorly controlled informal studies that lacked commonly recognized approaches for controlling for confounding factors. These may also have reflected early versions of the chair, since this writer has not found any more recent reports of discomfort at the knees in the last two decades.

Stranden (2000) pointed to the absence of evidence linking sitting on the Balans seat with loads on the knees, and described a prior unpublished study (Stranden, 1981) that also found no such evidence.

Furthermore, the flexion of the knee joint during upward seat deflection does not represent a likely cause for venous compression. This is evident from earlier studies on 'Balans' chairs, where venous pressure recordings at very much larger flexion did not indicate any venous obstruction at all (Stranden 1981). The most extreme sitting posture was applied at 'Balans Skulptur', where flexion was about 160 ± 1708 ; still with no venous obstruction affecting venous pressure profiles.

This writer has also reviewed partial reports from an in-house research study performed over nine months by physiotherapists at the Hospital of Robslend, Norway. They reported improvements in back pain but also acknowledged a potential for knee discomfort – which they said could be avoided through simple design changes.

Drury and Francher (1985) performed a small pilot study of 12 subjects sitting in Balans and conventional chairs for 2½ hours. Balans chair users reported higher rates of body part discomforts at the knees and (to a lesser extent). Even so, the authors noted, “a minority of subjects preferred this [Balans] chair to their own”.




Figure 4. The Thatsit Balans design enables users to promote circulation and reduce leg swelling by activating the venous pumping action of the legs, such as by rocking with one leg on the floor and the other leg on the shin support.

Stranden (2000) describes how the use of the three venous pumping systems synchronizes to aid circulation to the heart and prevent edema.

Dr. Grandjean's writings on the topic:

Several of Grandjean's editions of "Fitting the task to the man" (e.g., 1988 – see also Kroemer and Grandjean (1997)) reference an unpublished in-house report from the Swiss Federal Institute that was reputedly written in German by Krueger (1984), noting in each case.

"Krueger (1984) tested four models and found that the load on knees and lower legs is too high and sitting becomes painful after a while. (Some subjects even refused to sit longer than 2 h)."

Some others followed suit, describing this study – and one cannot help but wonder if it might have been taken as a secondary source from Grandjean without referencing it as such or reading the original material. Notably, Hermanau (1995, 1999) writes

"Studies by Krueger (1984), however, found that the load on the knees and lower legs is too great and sitting becomes painful."

Yet *The Swiss Federal Institute* indicated that they had no record of having sponsored such an unpublished German report and that it apparently not available directly from their Institute, leaving one to wonder whether recent articles that cite this report might be making inferences from secondary references to text that they have not actually read.

Through extensive follow through, Mr. Ed Miano of VarierUSA was able to locate Dr. Kruger (the correct spelling of his name), who had long since retired, but who graciously digitized and provided a copy of his original paper as a pdf. Dr. Kruger expressed surprise that his in-house review had attained so much notoriety, given the rather informal intent of the small review, which used 4 subjects on a short-term basis and was based on cursory reviews of subject's comments in the absence of objective data.

Both the translated paper provided by Mr. Miano and a Google-translation of the German version of the original paper each suggest that the findings misrepresented some important issues and content relevant to the paper.

The author noted that the higher seat height of the Balans was also rated positively for all users. He also wrote that all four of the subjects experienced some degree of discomfort the different versions of Balans chairs, but went on to describe the considerable variability in ratings between models. The chair that received the most positive ratings was the Balans Variable, which received positive scores by all users and "was found pleasant by all users".²⁷ Dr. Kruger noted that users continued to sit on this chair, without knee supports after the seating trials.

²⁷ One question that was left unanswered was whether the order of the chairs were counter balanced and otherwise controlled to avoid an order effect in testing. The Variable was listed last.

Conclusion

The research indicates that standing, sitting and semi-standing positions each have their benefits but also the limitations.

While many consider the spinal contours associated with the standing posture desirable, in research also tells us that in the real world many or most people do not want to stand all day – and when they work standing, often they assume less than ideal ergonomic positions.

We know that most users are accustomed and seem to prefer conventional seats with backrests and arm supports. These can provide important benefits by enabling some users to assume more neutral postures while transferring the loads from gravity to the backrests.

Yet many workers sitting in conventional chairs also spend much of the day with their back unsupported and their chair improperly adjusted – or not adjusted at all. As the day progresses, working postures often tend toward forward leaning anterior postures that put them at particular risk of discomfort and injuries to their spine and musculoskeletal system.

Balans chairs enable users to assume semi-standing postures. Users can slump in these chairs as well as conventional seating. Even so, research demonstrates the Balans seat can in some circumstances more effectively promote lumbar lordosis and though the debate rages on may eventually be found to be even “more neutral” than standing positions.

Almost all of the studies that assessed the Balans chair used versions of this chair that lacked back support. Yet such back supports provide important benefits, even if only to intermittently stretch and stabilize the spine. It seems reasonable to presume that the findings associated with the Balans would have been even stronger had these studies included back supports on the Balans chairs.

Further, while these studies compared the effects of sitting on conventional and Balans seats, there has been a lack of attention to the potential benefits for some users when alternating between both kinds of chairs. Yet for some users – particularly those who spend much of the time in forward oriented postures, the opportunity to alternate between these postures may provide important benefits.



References

- Aaras, A, Horgen, G. and Ro, O. (2000) Work with Visual Display Unit: Health consequences. International J Human-Computer Interaction. 12(1) 107-134.
- Adams, M. A., Green, T. P. and Dolan, P. (1994) The strength in anterior bending of lumbar intervertebral discs. Spine (Phila Pa 1976). 19, 19, 2197-2203.
- Adams, M.A., Hutton, W.C. (1983) The effect of posture on the fluid content of lumbar intervertebral discs. Spine. 8(6), 665-671.
- Adams, M. A., McMillan, D. W., Green, T. P. and Dolan, P. (1996a) Sustained loading generates stress concentrations in lumbar intervertebral discs. Spine (Phila Pa 1976). 21, 4, 434-438.
- Adams, M. A., McNally, D. S. and Dolan, P. (1996b) 'Stress' distributions inside intervertebral discs. The effects of age and degeneration. J Bone Joint Surg Br. 78, 6, 965-972.
- Adams, W., Stentz, T. L., Stonecipher, B. L. and Hallbeck, M.S. (1990) Comparison of spinal profiles while standing, supine, prone, and seated in four chair types: A pilot study. Human Factors and Ergonomics Society Annual Meeting Proceedings. 34, 679-683.
- Andersson, B. J., Ortengren, R., Nachemson, A. L., Elfstrom, G. and Broman, H. (1975) The sitting posture: an electromyographic and discometric study, Orthop Clin North Am. 6(1), 105-120.
- Andersson, B.J.G. and Ortengren, R. (1974) Lumbar disc pressure and myoelectric back muscle activity during sitting, Scandinavian Journal of Rehabilitation Medicine. 6, 115-121.
- Andersson, B.J.G., Ortengren, R., Nachemson, A. and Elfstrom, G. (1974) Lumbar disc pressure and myoelectric back muscle activity during sitting. Scandinavian J Rehabilitation Medicine. 6, 104-114.
- Andersson, G.B.J. (1980) The load on the lumbar spine in sitting postures. In: D.J. Osborne and J.A. Levis (Eds.) Human Factors in Transport Research. New York: Academic Press, 1980, 231-239.
- Andersson, G.B.J. (1986) Loads on the Spine during Sitting (Chapter 27). In: Corlett, E.N., and Eklund, J.A.E. and Manenica, I. (Eds.) The Ergonomics of Working Postures. Taylor & Francis: London. 309-318.
- Andersson, G.B.J. (1986) Loads on the spine during sitting. In: Corlett, N., Wilson, J. and Manenica, I. (Eds.) The Ergonomics of Working Postures. Philadelphia: Taylor and Francis. 109-118.
- Ankrum, D.R. (2000a) On the confusion between static load level and static task. Applied Ergonomics. 31(5), 545-546.

Ankrum, D.R. (2000b) Questions to ask when interpreting surface electromyography (SEMG) research. Proceedings of the Human Factors and Ergonomics Society Annual Meeting. 44, 530-533.

Ankrum, D.R. and Nemeth, K.J. (2000) Head and neck posture at computer workstations: What's neutral. Proceedings of the Human Factors and Ergonomics Society Annual Meeting. 44, 565-568.

Au, G., Cook, J. and McGill, S.M. (2001) Spinal shrinkage during repetitive controlled torsional, flexion and lateral bend motion exertions. Ergonomics. 44(4), 373-381.

Bazzucchi, I., Marchetti, M., Rosponi, A., Fattorini, L., Castellano, V., Sbriccoli, P. and Felici, F. (2005) Differences in the force/endurance relationship between young and older men. Eur J Applied Physiol. 93, 4, 390-397.

Bendix, T., Biering-Sorensen, F. (1983) Posture of the trunk when sitting on forward inclining seats. Scand J Rehabil Med. 15(4), 197-203.

Beach, T. A., Parkinson, R. J., Stothart, J. P. and Callaghan, J. P. (2005) Effects of prolonged sitting on the passive flexion stiffness of the in vivo lumbar spine. Spine J. 5, 2, 145-154.

Bendix, A., Jensen, C. and Bendix, T. (1988a) Posture, acceptability and energy consumption on a tiltable and a knee-support chair. Clinical Biomechanics. 3, 2, 66-73.

Bendix, T., Jessen, F. and Krohn, L. (1988b) Biomechanics of forward-reaching movements while sitting on fixed forward- or backward-inclining or tiltable seats. Spine (Phila Pa 1976). 13, 2, 193-196.

Bendix, T., Poulsen, V., Klausen, K. and Jensen, C. V. (1996) What does a backrest actually do to the lumbar spine? Ergonomics. 39, 4, 533-542.

Bennett, D. L., Gillis, D. K., Portney, L. G., Romanow, M., Sanchez, A. S. (1989) Comparison of integrated electromyographic activity and lumbar curvature during standing and during sitting in three chairs. Phys Ther. 69, 11, 902-913.

Bettany-Saltikov, J., Warren, J., Jobson, M. (2008) Ergonomically designed kneeling chairs are they worth it? : Comparison of sagittal lumbar curvature in two different seating postures. Stud Health Technol Inform. 140, 103-106.

Bhatnager, V., Drury, C. G., Schiro, S. G. (1985) Posture, postural discomfort, and performance. Human Factors. 27, 2, 189-199.

Bishu, R. R., Hallbeck, M.S., Riley, M. W. and Stentz, Terry L. (1991) Seating comfort and its relationship to spinal profile: A pilot study. International Journal of Industrial Ergonomics. 8, 1, 89-101.

Bridger, R. S. (1988) Postural adaptations to a sloping chair and work surface. Human Factors. 30, 2, 237-247.

Bridger, R. S., Wilkinson, D. and van Houweninge, T. (1989) Hip joint mobility and spinal angles in standing and in different sitting postures. Human Factors. 31, 2, 229-241.

Bridger, R.S., Kloote, C., Rowlands, B. and Fourie, G. (2000) Palliative interventions for sedentary low back pain: The physiotherapy ball, the kneeling chair and conventional ergonomics compared. Human Factors and Ergonomics Society Annual Meeting Proceedings. 44, 87-90.

Bridger, R.S., Von Eisenhart-Rothe, C. and Henneberg, M. (1989) Effects of Seat Slope and Hip Flexion on Spinal Angles in Sitting. Human Factors. 31, 679-688.

Brodeur, R.R. and Reynolds, H.M. (1990) Passive mechanics of the lumbo-pelvic spine for erect and slumped seated postures. Proceedings of the May 1990 International Conference on Spinal Manipulation. FCER: Arlington, Virginia. 190-193.

Brunswic, M. (1984a) Ergonomics of seat design. Physiotherapy. 70, 2, 40-43.

Brunswic, M., (1984b) Seat design in unsupported sitting. Proceedings of the 1984 Intl. Conf. Occup Ergonomics, Toronto 294-298.

Chaitow, L. (2009) Ligaments and positional release techniques? J Bodyw Mov Ther. 13, 2, 115-116.

Claus, A. P., Hides, J. A., Moseley, G. L. and Hodges, P. W. (2009) Is 'ideal' sitting posture real? Measurement of spinal curves in four sitting postures. Man Ther. 14, 4, 404-408.

Corlett, E.N. (2002) Personal communication.

Corlett, E.N. (1999) Are you sitting comfortably? International J Industrial Ergonomics, 24, 7-12.

Corlett, E.N. and Eklund, J.A. (1984) How does a backrest work? Applied Ergonomics. 15(2), 111-114.

Cram, J.R. and Vinitzky, I. (1995) Effects of chair design on back muscle fatigue. Journal of Occupational Rehabilitation. 5(2), 101-113.

De Carvalho, D.E., Dunk, N.M. and Callaghan, J.P. (2007) Gender differences in spinal posture and user positioning on a prototype seat pan. Proceedings of the American Biomechanics Society, 2007.

De Carvalho, D. E., Soave, D., Ross, K. and Callaghan, J. P. (2010) Lumbar spine and pelvic posture between standing and sitting: a radiologic investigation including reliability and repeatability of the lumbar lordosis measure. J Manipulative Physiol Ther. 33, 1, 48-55.

Dionne, C.E., Dunn, K.M. and Croft, P.R. (2006) Does back pain prevalence really decrease with increasing age? A systematic review. Age Ageing. 35(3), 229-234.

Dolan, P. and Adams, M. A. (2001) Recent advances in lumbar spinal mechanics and their significance for modeling. Clinical Biomechanics. 16, Supplement 1, S8-S16.

Dolan, K. J. and Green, A. (2006) Lumbar spine reposition sense: the effect of a 'slouched' posture. Man Ther. 11, 3, 202-207.

Dowell, W. R., Yuan, F. and Green, B. H. (2001) Office Seating Behaviors An Investigation of Posture, Task, and Job Type. Human Factors and Ergonomics Society Annual Meeting Proceedings. 45, 1245-1248.

Drury, C.G. and Francher, M. (1985) Evaluation of a forward-sloping chair. Applied Ergonomics. 16(1), 41-47.

Duncan, J. and Ferguson, D. (1974) Keyboard Operating Posture and Symptoms in Operating. Ergonomics, 17(5), 651-662.

Dunk, N. M. and Callaghan, J. P. (2005) Gender-based differences in postural responses to seated exposures. Clin Biomech (Bristol, Avon). 20, 10, 1101-1110.

Eklund, M. (1967) Prevalence of musculoskeletal disorders in office work. Socialmedicinsk, 6, 328-336.

Ericson, M.O. and Goldie, I. (1989) Spinal shrinkage with three different types of chair whilst performing video display unit work. International Journal of Industrial Ergonomics. 3(3), 177-183.

Evjenth, O. and Hamberg, J. (1985) Muscle Stretching in Manual Therapy. A Clinical Manual. Scand Book AB Sweden.

Fenety, P.A., Putnam, C. and Walker, J.M. (2000) In chair movement: Validity, reliability and implications for measuring sitting comfort. Applied Ergonomics. 31(4), 383-393.

Ferguson, S.A., Marras, W.S. and Gupta, P. (2000) Longitudinal quantitative measures of the natural course of low back pain recovery. Spine (Phila Pa 1976). 25(15), 1950-1956.

Fleischer, A.G., Rademacher, U. and Windberg, H.J. (1987) Individual characteristics of sitting behavior. Ergonomics. 30(4), 703-7.

Frey, J. K. and Tecklin, J. S. (1986) Comparison of lumbar curves when sitting on the Westnofa Balans Multi-Chair, sitting on a conventional chair, and standing. Phys Ther. 66, 9, 1365-1369.

Grandjean, E. (1976) Ergonomics of the Home. London: Taylor and Francis.

Grandjean, E., Hunting, W. and Pidermann, M. (1983) VDT workstation design: preferred settings and their effects. Human Factors. 25(2), 161-175.

Graf, M., Guggenbuhl, U. and Krueger, H. (1993) Investigations on the effects of seat shape and slope on posture, comfort and back muscle activity. International J Industrial Ergonomics, 12(1-2), 91-103.

Graf, M., Guggenbuhl, U. and Krueger, H. (1995) An assessment of seated activity and postures at five workplaces. International J Industrial Ergonomics, 15(2), 81-90.

Gregory, D. E., Dunk, N. M. and Callaghan, J. P. (2006) Stability ball versus office chair: comparison of muscle activation and lumbar spine posture during prolonged sitting. Hum Factors. 48, 1, 142-153.

Hermenau, D.C. (1999) Seating. Edited by Jacobs, K. Ergonomics for therapists. Woburn, MA: Butterworth-Heinemann. Citation on Page 203. Chapter 10, 219-237.

Hermenau, D.C. (1995) Seating. Edited by Jacobs, K. Ergonomics for therapists. Woburn, MA: Butterworth-Heinemann. Chapter 9.

Hunting, W., Grandjean, E. and Maeda, K. (1980) Constrained postures in accounting machine operators. Applied Ergonomics. 11(3), 145-149.

Hunting, W., Laubli, T. and Grandjean, E. (1981) Postural and visual loads at VDT workplaces: Constrained postures. Ergonomics. 24 (12), 917-931.

Hult, L. (1954) Cervical, dorsal and lumbar spine syndromes. Acta Orthopaedic Scandinavia (Supplement 17).

Keegan, J.J. (1953) Alterations of the lumbar curve related to posture and seating, J Bone and Joint Surgery. 35A (3), 589-603.

Kilbom, A. (1987). Short- and long-term effects of extreme physical inactivity: a brief review. In: Knave, B. and Wideback, P.G. (Eds.) Work with display units, Elsevier Science Publishers B V (North-Holland), Amsterdam, 219-228.

Kroemer, K. H. E. and Grandjean, E. (1997) Fitting the task to the human: a textbook of occupational ergonomics. New York: CRC Press.

Krueger H (1984). Zur Ergonomie von Balans-Sitzelementen im Hinblick auf ihre Verwendbarkeit als regulare Arbeitsstuhle. Report 8092. Zurich: Department of Ergonomics, Swiss Federal Institute of Technology.

Kumar, S. (2001) Theories of musculoskeletal injury causation. Ergonomics, 44, 1, 17-47.

Kumar, S., Narayan, Y. and Zedka, M. (1998) Strength in combined motions of rotation and flexion/ extension in normal young adults. Ergonomics 41, 1, 835-852.

Kumar, S. (2004) Ergonomics and biology of spinal rotation. Ergonomics. 47(4), 370-415.

Kumar, S., Narayan and Y. Zedka, M. (1996) An electromyographic study of unresisted trunk rotation with normal velocity among healthy subjects. Spine. 21(13), 1500-1512.

Lander, C., Korbon, G. A., DeGood, D. E., Rowlingson, J. C. (1987) The Balans chair and its semi-kneeling position: an ergonomic comparison with the conventional sitting position. Spine (Phila Pa 1976). 12, 3, 269-272.

Langdon, F.J. (1965). The design of cardpunches and the seating of operators. Ergonomics, 8, 61-68.

Le, B., Davidson, B., Solomonow, D., Zhou, B. H., Lu, Y., Patel, V. and Solomonow, M. (2009) Neuromuscular control of lumbar instability following static work of various loads. Muscle Nerve. 39, 1, 71-82.

Lengsfeld, M., Frank, A., van Deursen, D. L. and Griss, P. (2000a) Lumbar spine curvature during office chair sitting. Medical Engineering and Physics. 22(9), 665-669.

Lengsfeld, M., van Deursen, D. L., Rohlmann, A., van Deursen, L.L. and Griss, P. (2000b) Spinal load changes during rotatory dynamic sitting. Clinical Biomechanics (Bristol, Avon). 15(4), 295-297.

Leivseth, G. and Drerup, B. (1997) Spinal shrinkage during work in a sitting posture compared to work in a standing posture. Clinical Biomechanics. 12(7/8), 409-418.

Link, C. S., Nicholson, G. G., Shaddeau, S. A., Birch, R. and Gossman, M. R. (1990) Lumbar curvature in standing and sitting in two types of chairs: relationship of hamstring and hip flexor muscle length. Phys Ther. 70, 10, 611-618.

Lueder, R. (1994) Adjustability in context. Lueder, R. and Noro, K. (Ed): Hard Facts about Soft Machines, The Science of Seating, Philadelphia and London. Taylor and Francis. 25-35.

Lueder, R. (2005) Ergonomics of sitting and seating: The case for and against movement for its own sake. An ergonomics review of the literature. Sponsored by Allsteel Seating. August 4, 2004. Updated June 5, 2005. 30 pp. www.humanics-es.com/movement-ergonomics.htm

Lueder, R. (2008) Physical development in children and adolescents and age-related risks. In: Lueder, R. and Rice, V.B. (Eds.) Ergonomics for Children: Designing products and places for toddlers to teens. London: Taylor & Francis.

Mandal, A.C. (1976) Work-chair with tilting seat. Ergonomics. 19(2), 157-164.

Mandal, A.C. (1981) The seated man (homo sedens) the seated work position. Theory and practice. Applied Ergonomics. 12(1), 19-26.

Mandal, A.C. (1982) The seated man. Theories and realities. Human Factors and Ergonomics Society Annual Meeting Proceedings. 26, 520-524.

Mariconda, M., Galasso, O., Imbimbo, L., Lotti, G. and Milano, C. (2007) Relationship between alterations of the lumbar spine, visualized with magnetic resonance imaging, and occupational variables. Eur Spine J. 16(2), 255-266.

NIOSH (1997) Low Back Musculoskeletal Disorders: Evidence for Work-Relatedness (Chapter 6). In: Bernard, B.P. (Ed.) Musculoskeletal Disorders (MSDs) and Workplace Factors. A Critical Review of Epidemiologic Evidence for Work-Related Musculoskeletal Disorders of the Neck, Upper Extremity, and Low Back. National Dept. Health & Human Services. National Institute for Occupational Safety & Health. DHHS. (NIOSH) Publication No. 97-141 (1997). Second printing.

O'Sullivan, P. B., Dankaerts, W., Burnett, A. F., Farrell, G. T., Jefford, E., Naylor, C. S. and O'Sullivan, K. J. (2006) Effect of different upright sitting postures on spinal-pelvic curvature and trunk muscle activation in a pain-free population. Spine (Phila Pa 1976). 31(19), E707-712.

Panjabi, M., Yamamoto, I., Oxland, T. and Crisco, J. (1989) How does posture affect coupling in the lumbar spine? Spine (Phila Pa 1976). 14, 9, 1002-1011.

Panjabi, M. and White, A. (2001) Biomechanics in the Musculoskeletal System. Churchill Livingstone.

Pynt, J. (2010) Ancient Greek Seating: A cause of modern day back pain. Intervention & Prevention in Health: Research to Practice. Pynt, J. (Ed.) Antarctica StuVac Seminar Proceedings. Antarctica, February 2010. 1-32.

Pynt, J. and Higgs, J. (2010. in Press). Healthy Seating: A History. NY: Cambria Press. In Press.

Pynt, J., Higgs, J. and Mackey, M. (2001) Seeking the optimal posture of the seated lumbar spine. Physiotherapy Theory and Practice. 17. 5-21.

Pynt, J., Higgs, J. and Mackey, M. (2002) Historical perspective milestones in the evolution of lumbar spinal postural health in seating. Spine (Phila Pa 1976). 27, 19, 2180-2189.

Pynt, J., Mackey, M. G. and Higgs, J. (2008) Kyphosed seated postures: extending concepts of postural health beyond the office. J Occup Rehabil. 18, 1, 35-45.

Ramazzini (1777) Essai sur les Maladies de Disseus. Original translation from Latin in "De Mortis Artificum" by M. De Foureau.

Rebiffe, R. (1980). General reflections on the postural comfort of the driver and passengers: Consequences on seat design. In D.J. Osborne and J.A. Levis (eds.), Human Factors in Transport Research. Volume 2 - User Factors: Comfort, the Environment and Behaviour, (pp. 240-248). New York: Academic Press.

Reinecke, S., Weisman, G. and Pope, M.H. (1994b) Effects of body position and center of gravity on tolerance of seated postures. Lueder, R. and Noro, K. (Ed): Hard Facts about Soft Machines. The Science of Seating. Philadelphia and London. Taylor and Francis. 165-171.

Rogers, E. L. and Granata, K. P. (2006) Disturbed paraspinal reflex following prolonged flexion-relaxation and recovery. Spine (Phila Pa 1976). 31, 7, 839-845.

Rohlmann, A., Arntz, U., Graichen, F., Bergmann, G. (2001) Loads on an internal spinal fixation device during sitting. J Biomechanics. 34, 8, 989-993.

Rohlmann, A., Neller, S., Claes, L., Bergmann, G., Wilke, H. J. (2001) Influence of a follower load on intradiscal pressure and intersegmental rotation of the lumbar spine. Spine (Phila Pa 1976). 26, 24, E557-561.

Shenoy, S., Aruin, A. S. (2007) Effect of chair design on feed-forward postural control in sitting. Motor Control. 11, 4, 309-321.

Smoliga, J. M., Myers, J. B., Redfern, M. S., Lephart, S. M. (2010) Reliability and precision of EMG in leg, torso, and arm muscles during running. J Electromyogr Kinesiol. 20, 1, e1-9.

Solomonow, M. (2009) Ligaments: a source of musculoskeletal disorders. J Bodywork Movement Therapy. 13, 2, 136-154.

Solomonow, M., Baratta, R. V., Banks, A., Freudenberger, C., Zhou, B. H. (2003a) Flexion-relaxation response to static lumbar flexion in males and females. Clin Biomech (Bristol, Avon). 18, 4, 273-279.

Straker, L.M., O'Sullivan, P.B., Smith, A.J., Perry, M.C., Coleman, J. (2008) Sitting spinal posture in adolescents differs between genders, but is not clearly related to neck/shoulder pain: An observational study. Aust J Physiother. 54(2), 127-133.

Stranden, E. (1981) The influence of 'Balans' chair sitting on ankle vein pressure. Unpublished exhibition leaflet. International Design Exhibition, New York, May. Cited in Stranden (2000, op cit).

Stranden, E. (2000) Dynamic leg volume changes when sitting in a locked and free floating tilt office chair. Ergonomics. 43, 3, 421-433.

Van Deursen, D.L., Snijders, C.J., Van Dieen, J.H., Kingma, I., Van Deursen, L.L. (2001) The effect of passive vertebral rotation on pressure in the nucleus pulposus. J Biomech. 34(3), 405-408.

Vink, P., Porcar-Seder, R., De, P., Lvaro, P., Krause, F. (2007) Office chairs are often not adjusted by end-users. Human Factors and Ergonomics Society Annual Meeting Proceedings. 51, 1015-1019.

Vink, P., Konijn, I., Jongejan, B., Berger, M., (2009) Varying the office work posture between standing, half-standing and sitting results in less discomfort. Proceedings of the International Conference on Ergonomics and Health Aspects of Work with Computers: Held as Part of HCI International 2009. San Diego, CA Springer-Verlag 115-120

Wilke, H. J., Neef, P., Caimi, M., Hoogland, T., Claes, L. E. (1999) New in vivo measurements of pressures in the intervertebral disc in daily life. Spine. 24, 8, 755-762.

Yamaguchi, Y. and Umezawa, F. (1973) Development of a chair to minimize disc distortion in the sitting position. Unpublished presentation at the 4th International Congress on Ergonomics. Strasbourg, July 1970. Cited in: Grandjean, E. (1976) Ergonomics of the Home. London: Taylor and Francis.

Additional Research

APHA (2003) Deep-vein thrombosis: Advancing awareness to protect patient lives. American Public Health Association. Available at www.apha.org/NR/rdonlyres/A209F84A-7C0E-4761-9ECF-61D22E1E11F7/0/DVT_White_Paper.pdf

Astfalck, R. G., O'Sullivan P, B., Straker, L. M., Smith, A. J., Burnett, A., Caneiro, J. P., Dankaerts, W. (2010) Sitting Postures and Trunk Muscle Activity in Adolescents With and Without Nonspecific Chronic Low Back Pain: An Analysis Based on Subclassification. Spine (Phila Pa 1976).

Bjorck-van Dijken, C., Fjellman-Wiklund, A., Hildingsson, C. (2008) Low back pain, lifestyle factors and physical activity: a population based-study. J Rehabil Med. 40, 10, 864-869.

Blouin, J. S., Descarreaux, M., Belanger-Gravel, A., Simoneau, M., Teasdale, N. (2003) Attenuation of human neck muscle activity following repeated imposed trunk-forward linear acceleration. Exp Brain Res. 150, 4, 458-464.

Brand, J. L. (2008) Office Ergonomics: A Review of Pertinent Research and Recent Developments. Reviews of Human Factors and Ergonomics. 4, 245-282.

Bridger, R.S., Kloote, C., Rowlands, B., Fourie, G. (2000) Palliative interventions for sedentary low back pain: The physiotherapy ball, the kneeling chair and conventional ergonomics compared. Human Factors and Ergonomics Society Annual Meeting Proceedings. 44, 87-90.

Brown, T. N., Palmieri-Smith, R. M., McLean, S. G. (2009) Sex and limb differences in hip and knee kinematics and kinetics during anticipated and unanticipated jump landings: implications for anterior cruciate ligament injury. British Journal of Sports Medicine. 43, 13, 1049-1056.

Burnett, A., O'Sullivan, P., Ankarberg, L., Gooding, M., Nelis, R., Offermann, F., Persson, J. (2008) Lower lumbar spine axial rotation is reduced in end-range sagittal postures when compared to a neutral spine posture. Man Ther. 13, 4, 300-306.

Burton, A. K. (1984) Electromyography and office-chair design, a pilot study. Behaviour & Information Technology. 3, 4, 353 - 357.

Callaghan, J. P., Dunk, N. M. (2002) Examination of the flexion relaxation phenomenon in erector spinae muscles during short duration slumped sitting. Clin Biomech (Bristol, Avon). 17, 5, 353-360.

Callaghan, J. P., McGill, S. M. (2001) Low back joint loading and kinematics during standing and unsupported sitting. Ergonomics. 44, 3, 280-294.

Carcone, S.M., Keir, P.J. (2007) Effects of backrest design on biomechanics and comfort during seated work. Applied Ergon. 38(6), 755-764.

Carpes, F. P., Dagnese, F., Kleinpaul, J. F., Martins Ede, A., Mota, C. B. (2009) Effects of workload on seat pressure while cycling with two different saddles. J Sex Med. 6, 10, 2728-2735.

Chany, A. M., Parakkat, J., Yang, G., Burr, D. L., Marras, W. S. (2006) Changes in spine loading patterns throughout the workday as a function of experience, lift frequency, and personality. Spine J. 6, 3, 296-305.

Christie, H. J., Kumar, S., Warren, S. A. (1995) Postural aberrations in low back pain. Arch Phys Med Rehabil. 76, 3, 218-224.

Chu, D., LeBlanc, R., D'Ambrosia, P., D'Ambrosia, R., Baratta, R. V., Solomonow, M. (2003) Neuromuscular disorder in response to anterior cruciate ligament creep. Clin Biomech (Bristol, Avon). 18, 3, 222-230.

Cramer, G. D., Cantu, J. A., Dorsett, R. D., Greenstein, J. S., McGregor, M., Howe, J. E., Glenn, W. V. (2003) Dimensions of the lumbar intervertebral foramina as determined from the sagittal plane magnetic resonance imaging scans of 95 normal subjects. J Manipulative Physiol Ther. 26, 3, 160-170.

Cramer, G. D., Cantu, J. A., Pocius, J. D., Cambron, J. A., McKinnis, R. A. (2010) Reliability of zygapophysial joint space measurements made from magnetic resonance imaging scans of acute low back pain subjects: comparison of 2 statistical methods. J Manipulative Physiol Ther. 33, 3, 220-225.

Cramer, G. D., Gregerson, D. M., Knudsen, J. T., Hubbard, B. B., Ustas, L. M., Cantu, J. A. (2002) The effects of side-posture positioning and spinal adjusting on the lumbar Z joints: a randomized controlled trial with sixty-four subjects. Spine (Phila Pa 1976). 27, 22, 2459-2466.

Cramer, G. D., Scott, C. Y., Tuck, N. R. (1998) The holey spine: a summary of the history of scientific investigation of the intervertebral foramina. Chiropr Hist. 18, 2, 13-24.

Cramer, G. D., Skogsbergh, D. R., Bakkum, B. W., Winterstein, J. F., Yu, S., Tuck, N. R., Jr. (2002) Evaluation of transforaminal ligaments by magnetic resonance imaging. J Manipulative Physiol Ther. 25, 4, 199-208.

Cramer, G. D., Tuck, N. R., Jr., Knudsen, J. T., Fonda, S. D., Schliesser, J. S., Fournier, J. T., Patel, P. (2000) Effects of side-posture positioning and side-posture adjusting on the lumbar zygapophysial joints as evaluated by magnetic resonance imaging: a before and after study with randomization. J Manipulative Physiol Ther. 23, 6, 380-394.

Dankaerts, W., O'Sullivan, P. B., Burnett, A. F., Straker, L. M. (2007) The use of a mechanism-based classification system to evaluate and direct management of a patient with non-specific chronic low back pain and motor control impairment--a case report. Man Ther. 12, 2, 181-191.

Dankaerts, W., O'Sullivan, P. B., Straker, L. M., Burnett, A. F., Skouen, J. S. (2006) The inter-examiner reliability of a classification method for non-specific chronic low back pain patients with motor control impairment. Man Ther. 11, 1, 28-39.

Dankaerts, W., O'Sullivan, P., Burnett, A., Straker, L. (2006) Altered patterns of superficial trunk muscle activation during sitting in nonspecific chronic low back pain patients: importance of subclassification. Spine (Phila Pa 1976). 31, 17, 2017-2023.

Dankaerts, W., O'Sullivan, P., Burnett, A., Straker, L. (2006) Differences in sitting postures are associated with nonspecific chronic low back pain disorders when patients are subclassified. Spine (Phila Pa 1976). 31, 6, 698-704.

Dankaerts, W., O'Sullivan, P., Burnett, A., Straker, L., Davey, P., Gupta, R. (2009) Discriminating healthy controls and two clinical subgroups of nonspecific chronic low back pain patients using trunk muscle activation and lumbosacral kinematics of postures and movements: a statistical classification model. Spine (Phila Pa 1976). 34, 15, 1610-1618.

de Carvalho, D.E., Dunk, N.M., Callaghan, J.P. (2007) Gender differences in spinal posture and user positioning in a prototype seat pan. Annual Meeting of the American Society of Biomechanics. Palo Alto 2007.

de Looze, M. P., Kuijt-Evers, L. F., van Dieen, J. (2003) Sitting comfort and discomfort and the relationships with objective measures. Ergonomics. 46, 10, 985-997.

Delleman, Nico J. (2000) Evaluation of Head and Neck Postures. Human Factors and Ergonomics Society Annual Meeting Proceedings. 44, 732-735.

Demaree, H. A., Higgins, D. A., Williamson, J., Harrison, D. W. (2002) Asymmetry in hand grip strength and fatigue in low- and high-hostile men. Int J Neurosci. 112, 4, 415-428.

Descarreaux, M., Blouin, J. S., Teasdale, N. (2005) Isometric force production parameters during normal and experimental low back pain conditions. BMC Musculoskeletal Disorders. 6, 6.

Descarreaux, M., Lafond, D., Jeffrey-Gauthier, R., Centomo, H., Cantin, V. (2008) Changes in the flexion relaxation response induced by lumbar muscle fatigue. BMC Musculoskeletal Disorders. 9, 10.

Deyo, R. A., Mirza, S. K., Martin, B. I. (2006) Back pain prevalence and visit rates: estimates from U.S. national surveys, 2002. Spine (Phila Pa 1976). 31, 23, 2724-2727.

Deyo, R. A., Mirza, S. K., Turner, J. A., Martin, B. I. (2009) Overtreating chronic back pain: time to back off? J Am Board Fam Med. 22, 1, 62-68.

Dionne, C.E., Dunn, K.M., Croft, P.R., Nachemson, A.L., Buchbinder, R., Walker, B.F., Wyatt, M., Cassidy, J.D., Rossignol, M., Leboeuf-Yde, C., Hartvigsen, J., Leino-Arjas, P., Latza, U., Reis, S., Gil Del Real, M.T., Kovacs, F.M., Oberg, B., Cedraschi, C., Bouter, L.M., Koes, B.W., Picavet, H.S., Van Tulder, M.W., Burton, K., Foster, N.E., Macfarlane, G.J., Thomas, E., Underwood, M., Waddell, G., Shekelle, P., Volinn, E., Von Korff, M. (2008) A consensus approach toward the standardization of back pain definitions for use in prevalence studies. Spine (Phila Pa 1976). 33(1), 95-103.

Drummond, D. S., Narechania, R. G., Rosenthal, A. N., Breed, A. L., Lange, T. A., Drummond, D. K. (1982) A study of pressure distributions measured during balanced and unbalanced sitting. J Bone Joint Surg [Am]. 64, 7, 1034-1039.

Drummond, D. S., Rogala, E., Templeton, J., Cruess, R. (1974) Proximal hamstring release for knee flexion and crouched posture in cerebral palsy. J Bone Joint Surg [Am]. 56, 8, 1598-1602.

Dunk, N. M., Kedgley, A. E., Jenkyn, T. R., Callaghan, J. P. (2009) Evidence of a pelvis-driven flexion pattern: are the joints of the lower lumbar spine fully flexed in seated postures? Clin Biomech (Bristol, Avon). 24, 2, 164-168.

Fennell, A. J., Jones, A. P., Hukins, D. W. (1996) Migration of the nucleus pulposus within the intervertebral disc during flexion and extension of the spine. Spine (Phila Pa 1976). 21, 23, 2753-2757.

Fersum, K. V., Dankaerts, W., O'Sullivan, P. B., Maes, J., Skouen, J. S., Bjordal, J. M., Kvale, A. (2009) Integration of sub-classification strategies in RCTs evaluating manual therapy treatment and exercise therapy for non-specific chronic low back pain (NSCLBP): a systematic review. Br J Sports Med.

Fjellman-Wiklund, A., Sundelin, G. (1998) Musculoskeletal discomfort of music teachers: an eight-year perspective and psychosocial work factors. Int J Occup Environ Health. 4, 2, 89-98.

Floyd, W. F., Silver, P. H. (1955) The function of the erector spinae muscles in certain movements and postures in man. J Physiol. 129, 1, 184-203.

Gallucci, R. M., Lee, E. G., Tomasek, J. J. (2006) IL-6 modulates alpha-smooth muscle actin expression in dermal fibroblasts from IL-6-deficient mice. J Invest Dermatol. 126, 3, 561-568.

Gerr, F., Marcus, M. (1999) Risk factors for carpal tunnel syndrome. Arch Intern Med. 159, 9, 1008-1010.

Gerr, F., Marcus, M. (2008) Yes, the "one-handed" scientist lacks rigor--why investigators should not use causal language when interpreting the results of a single study. Am J Ind Med. 51, 10, 795-796, author reply 797-798.

Gerr, F., Marcus, M., Ensor, C., Kleinbaum, D., Cohen, S., Edwards, A., Gentry, E., Ortiz, D. J., Monteilh, C. (2002) A prospective study of computer users: I. Study design and incidence of musculoskeletal symptoms and disorders. Am J Ind Med. 41, 4, 221-235.

Gerr, F., Marcus, M., Monteilh, C. (2004) Epidemiology of musculoskeletal disorders among computer users: lesson learned from the role of posture and keyboard use. J Electromyogr Kinesiol. 14, 1, 25-31.

Gerr, F., Marcus, M., Monteilh, C., Hannan, L., Ortiz, D., Kleinbaum, D. (2005) A randomised controlled trial of postural interventions for prevention of musculoskeletal symptoms among computer users. Occup Environ Med. 62, 7, 478-487.

Gerr, F., Marcus, M., Ortiz, D. J. (1996) Methodological limitations in the study of video display terminal use and upper extremity musculoskeletal disorders. Am J Ind Med. 29, 6, 649-656.

Gerr, F., Marcus, M., Ortiz, D., White, B., Jones, W., Cohen, S., Gentry, E., Edwards, A., Bauer, E. (2000) Computer users' postures and associations with workstation characteristics. AIHAJ. 61, 2, 223-230.

Gerr, F., Monteilh, C. P., Marcus, M. (2006) Keyboard use and musculoskeletal outcomes among computer users. J Occup Rehabil. 16, 3, 265-277.

Goossens, R. H., Snijders, C. J., Roelofs, G. Y., van Buchem, F. (2003) Free shoulder space requirements in the design of high backrests. Ergonomics. 46, 5, 518-530.

Granata, K. P., Rogers, E., Moorhouse, K. (2005) Effects of static flexion-relaxation on paraspinal reflex behavior. Clin Biomech (Bristol, Avon). 20, 1, 16-24.

Gregory, D. E., Dunk, N. M., Callaghan, J. P. (2006) Stability ball versus office chair: comparison of muscle activation and lumbar spine posture during prolonged sitting. Hum Factors. 48, 1, 142-153.

Hannan, L. M., Monteilh, C. P., Gerr, F., Kleinbaum, D. G., Marcus, M. (2005) Job strain and risk of musculoskeletal symptoms among a prospective cohort of occupational computer users. Scand J Work Environ Health. 31, 5, 375-386.

Hansen, F. R., Bendix, T., Skov, P., Jensen, C. V., Kristensen, J. H., Krohn, L., Schioeler, H. (1993) Intensive, dynamic back-muscle exercises, conventional physiotherapy, or placebo-control treatment of low-back pain. A randomized, observer-blind trial. Spine (Phila Pa 1976). 18, 1, 98-108.

Harkness, E. F., Macfarlane, G. J., Nahit, E. S., Silman, A. J., McBeth, J. (2003) Mechanical and psychosocial factors predict new onset shoulder pain: a prospective cohort study of newly employed workers. Occup Environ Med. 60, 11, 850-857.

Harkness, E. F., Macfarlane, G. J., Nahit, E. S., Silman, A. J., McBeth, J. (2003) Risk factors for new-onset low back pain amongst cohorts of newly employed workers. Rheumatology (Oxford). 42, 8, 959-968.

Harkness, E. F., Macfarlane, G. J., Nahit, E., Silman, A. J., McBeth, J. (2004) Mechanical injury and psychosocial factors in the work place predict the onset of widespread body pain: a two-year prospective study among cohorts of newly employed workers. Arthritis Rheum. 50, 5, 1655-1664.

Harkness, E. F., Nahit, E. S., Macfarlane, G. J., Silman, A. J., McBeth, J., Dunn, G. (2003) Generalised estimating equations and low back pain. Occup Environ Med. 60, 5, 378-380, author reply 380-371.

Harrison, D. D., Harrison, S. O., Croft, A. C., Harrison, D. E., Troyanovich, S. J. (1999) Sitting biomechanics part I: review of the literature. J Manipulative Physiol Ther. 22, 9, 594-609.

Harrison, D. D., Harrison, S. O., Croft, A. C., Harrison, D. E., Troyanovich, S. J. (2000) Sitting biomechanics, part II: optimal car driver's seat and optimal driver's spinal model. J Manipulative Physiol Ther. 23, 1, 37-47.

Hassaballa, M. A., Porteous, A. J., Learmonth, I. D. (2007) Functional outcomes after different types of knee arthroplasty: kneeling ability versus descending stairs. Med Sci Monit. 13, 2, CR77-81.

Hassaballa, M. A., Porteous, A. J., Newman, J. H. (2004) Observed kneeling ability after total, unicompartmental and patellofemoral knee arthroplasty: perception versus reality. Knee Surg Sports Traumatol Arthrosc. 12, 2, 136-139.

Hassaballa, M. A., Porteous, A. J., Newman, J. H., Rogers, C. A. (2003) Can knees kneel? Kneeling ability after total, unicompartmental and patellofemoral knee arthroplasty. Knee. 10, 2, 155-160.

Hedge, A., Jagdeo, J., Agarwal, A., Rockey-Harris, K. (2005) Sitting or standing for computer work does a negative-tilt keyboard tray make a difference. Human Factors and Ergonomics Society Annual Meeting Proceedings. 49, 808-812.

Hermans, V., Hautekiet, M., Haex, B., Spaepen, A. J., Van der Perre, G. (1999) Lipoatrophia semicircularis and the relation with office work. Applied Ergonomics. 30, 4, 319-324.

Hildebrandt, W., Herrmann, J., Stegemann, J. (1993) Vascular adjustment and fluid reabsorption in the human forearm during elevation. Eur J Applied Physiol Occup Physiol. 66, 5, 397-404.

Hildebrandt, W., Herrmann, J., Stegemann, J. (1994) Fluid balance versus blood flow autoregulation in the elevated human limb: the role of venous collapse. Eur J Appl Physiol Occup Physiol. 69, 2, 127-131.

Hinz, B., Celetta, G., Tomasek, J. J., Gabbiani, G., Chaponnier, C. (2001) Alpha-smooth muscle actin expression upregulates fibroblast contractile activity. Mol Biol Cell. 12, 9, 2730-2741.

Hoops, H., Zhou, B. H., Lu, Y., Solomonow, M., Patel, V. (2007) Short rest between cyclic flexion periods is a risk factor for a lumbar disorder. Clin Biomech (Bristol, Avon). 22, 7, 745-757.

Husemann, B., Von Mach, C.Y., Borsotto, D., Zepf, K.I., Scharnbacher, J. (2009) Comparisons of musculoskeletal complaints and data entry between a sitting and a sit-stand workstation paradigm. Hum Factors. 51(3), 310-320.

Ivancic, P. C., Coe, M. P., Ndu, A. B., Tominaga, Y., Carlson, E. J., Rubin, W., Dipl-Ing, F. H., Panjabi, M. M. (2007) Dynamic mechanical properties of intact human cervical spine ligaments. Spine J. 7(6), 659-665.

Janwantanakul, P., Pensri, P., Jiamjarasrangsi, V., Sinsongsook, T. (2008) Prevalence of self-reported musculoskeletal symptoms among office workers. Occup Med (Lond). 58, 6, 436-438.

Jenkins, C., Barker, K. L., Pandit, H., Dodd, C. A., Murray, D. W. (2008) After partial knee replacement, patients can kneel, but they need to be taught to do so: a single-blind randomized controlled trial. Phys Ther. 88, 9, 1012-1021.

Jensen, T. S., Kjaer, P., Korsholm, L., Bendix, T., Sorensen, J. S., Manniche, C., Leboeuf-Yde, C. (2009) Predictors of new vertebral endplate signal (Modic) changes in the general population. Eur Spine J.

Jones, G. T., Harkness, E. F., Nahit, E. S., McBeth, J., Silman, A. J., Macfarlane, G. J. (2007) Predicting the onset of knee pain: results from a 2-year prospective study of new workers. Ann Rheum Dis. 66, 3, 400-406.

Karacan, I., Aydin, T., Sahin, Z., Cidem, M., Koyuncu, H., Aktas, I., Uludag, M. (2004) Facet angles in lumbar disc herniation: their relation to anthropometric features. Spine (Phila Pa 1976). 29, 10, 1132-1136.

Kee, D. (2002) A method for analytically generating three-dimensional isocomfort workspace based on perceived discomfort. Applied Ergonomics. 33, 1, 51-62.

Kee, D., Karwowski, W. (2001) The boundaries for joint angles of isocomfort for sitting and standing males based on perceived comfort of static joint postures. Ergonomics. 44, 6, 614-648.

Kee, D., Karwowski, W. (2004) Joint angles of isocomfort for female subjects based on the psychophysical scaling of static standing postures. Ergonomics. 47, 4, 427-445.

Klausen, K., Rasmussen, B. (1968) On the location of the line of gravity in relation to L5 in standing. Acta Physiol Scand. 72, 1, 45-52.

Klussmann, A., Gebhardt, H., Liebers, F., Rieger, M. A. (2008) Musculoskeletal symptoms of the upper extremities and the neck: a cross-sectional study on prevalence and symptom-predicting factors at visual display terminal (VDT) workstations. BMC Musculoskelet Disord. 9, 96.

LaBry, R., Sbriccoli, P., Zhou, B. H., Solomonow, M. (2004) Longer static flexion duration elicits a neuromuscular disorder in the lumbar spine. J Applied Physiol. 96, 5, 2005-2015.

Le, B., Davidson, B., Solomonow, D., Zhou, B. H., Lu, Y., Patel, V., Solomonow, M. (2009) Neuromuscular control of lumbar instability following static work of various loads. Muscle Nerve. 39, 1, 71-82.

Leijnse, J. N., Bonte, J. E., Landsmeer, J. M., Kalker, J. J., Van der Meulen, J. C., Snijders, C. J. (1992) Biomechanics of the finger with anatomical restrictions--the significance for the exercising hand of the musician. J Biomech. 25, 11, 1253-1264.

Li, L., Patel, N., Solomonow, D., Le, P., Hoops, H., Gerhardt, D., Johnson, K., Zhou, B. H., Lu, Y., Solomonow, M. (2007) Neuromuscular response to cyclic lumbar twisting. Human Factors. 49, 5, 820-829.

Lin, F., Parthasarathy, S., Taylor, S. J., Pucci, D., Hendrix, R. W., Makhsous, M. (2006) Effect of different sitting postures on lung capacity, expiratory flow, and lumbar lordosis. Arch Phys Med Rehabil. 87, 4, 504-509.

Little, J. S., Khalsa, P. S. (2005) Human Lumbar Spine Creep during Cyclic and Static Flexion: Creep Rate, Biomechanics, and Facet Joint Capsule Strain. Annals of Biomedical Engineering. 33, 3, 391-401.

Lou, S. Z., Chou, Y. L., Chou, P. H., Lin, C. J., Chen, U. C., Su, F. C. (2001) Sit-to-stand at different periods of pregnancy. Clin Biomech (Bristol, Avon). 16, 3, 194-198.

Lu, D., Solomonow, M., Zhou, B., B., R. V., Li, L. (2004) Frequency-dependent changes in neuromuscular responses to cyclic lumbar flexion. Journal of Biomechanics. 37, 6, 845-855.

MacLean, J. J., Lee, C. R., Grad, S., Ito, K., Alini, M., Iatridis, J. C. (2003) Effects of immobilization and dynamic compression on intervertebral disc cell gene expression in vivo. Spine (Phila Pa 1976). 28, 10, 973-981.

Madsen, R., Jensen, T. S., Pope, M., Sorensen, J. S., Bendix, T. (2008) The effect of body position and axial load on spinal canal morphology: an MRI study of central spinal stenosis. Spine (Phila Pa 1976). 33, 1, 61-67.

Magnusson, M. L., Aleksiev, A. R., Spratt, K. F., Lakes, R. S., Pope, M. H. (1996) Hyperextension and spine height changes. Spine (Phila Pa 1976). 21, 22, 2670-2675.

Magnusson, M., Hansson, T., Pope, M. H. (1994) The effect of seat back inclination on spine height changes. Applied Ergonomics. 25, 5, 294-298.

Magnusson, M., Pope, M. (1996) Body height changes with hyperextension. Clin Biomech (Bristol, Avon). 11, 4, 236-238.

Magnusson, M., Pope, M. (1997) Body height changes with hyperextension. Clin Biomech (Bristol, Avon). 12, 2, 1.

Magnusson, M., Pope, M. H., Hansson, T. (1996) Does a back support have a positive biomechanical effect? Applied Ergonomics. 27, 3, 201-205.

Maigne, J. Y., Guillon, F. (2000) Highlighting of intervertebral movements and variations of intradiskal pressure during lumbar spine manipulation: a feasibility study. J Manipulative Physiol Ther. 23, 8, 531-535.

Maigne, J. Y., Lapeyre, E., Morvan, G., Chatellier, G. (2003) Pain immediately upon sitting down and relieved by standing up is often associated with radiologic lumbar instability or marked anterior loss of disc space. Spine (Phila Pa 1976). 28, 12, 1327-1334.

Majeske, C., Buchanan, C. (1984) Quantitative description of two sitting postures. With and without a lumbar support pillow. Phys Ther. 64, 10, 1531-1535.

Makhsous, M., Lin, F., Bankard, J., Hendrix, R. W., Hepler, M., Press, J. (2009) Biomechanical effects of sitting with adjustable ischial and lumbar support on occupational

low back pain: evaluation of sitting load and back muscle activity. BMC Musculoskelet Disord. 10, 17.

Makhsous, M., Lin, F., Hendrix, R. W., Hepler, M., Zhang, L. Q. (2003) Sitting with adjustable ischial and back supports: biomechanical changes. Spine (Phila Pa 1976). 28, 11, 1113-1121, discussion 1121-1112.

Marcus, M., Gerr, F. (1996) Upper extremity musculoskeletal symptoms among female office workers: associations with video display terminal use and occupational psychosocial stressors. Am J Ind Med. 29, 2, 161-170.

Marcus, M., Gerr, F., Monteilh, C., Ortiz, D. J., Gentry, E., Cohen, S., Edwards, A., Ensor, C., Kleinbaum, D. (2002) A prospective study of computer users: II. Postural risk factors for musculoskeletal symptoms and disorders. Am J Ind Med. 41, 4, 236-249.

Marras, W. S., Parakkat, J., Chany, A. M., Yang, G., Burr, D., Lavender, S. A. (2006) Spine loading as a function of lift frequency, exposure duration, and work experience. Clin Biomech (Bristol, Avon). 21, 4, 345-352.

Martin, B. I., Deyo, R. A., Mirza, S. K., Turner, J. A., Comstock, B. A., Hollingworth, W., Sullivan, S. D. (2008) Expenditures and health status among adults with back and neck problems. JAMA. 299, 6, 656-664.

Masharawi, Y. M., Kjaer, P., Bendix, T., Manniche, C., May, H., Mirovsky, Y., Anekshtein, Y., Jensen, T. S., Hershkovitz, I. (2009) Lumbar facet and interfacet shape variation during growth in children from the general population: a three-year follow-up MRI study. Spine (Phila Pa 1976). 34, 4, 408-412.

Maurer, C. L., Sprigle, S. (2004) Effect of seat inclination on seated pressures of individuals with spinal cord injury. Phys Ther. 84, 3, 255-261.

McBeth, J., Harkness, E. F., Silman, A. J., Macfarlane, G. J. (2003) The role of workplace low-level mechanical trauma, posture and environment in the onset of chronic widespread pain. Rheumatology (Oxford). 42, 12, 1486-1494.

McMillan, D. W., Garbutt, G., Adams, M. A. (1996) Effect of sustained loading on the water content of intervertebral discs: implications for disc metabolism. Ann Rheum Dis. 55, 12, 880-887.

Mitchell, T., O'Sullivan, P. B., Burnett, A. F., Straker, L., Smith, A. (2008) Regional differences in lumbar spinal posture and the influence of low back pain. BMC Musculoskeletal Disorders. 9, 152.

Moore, S., Brunt, D. (1991) Effects of trunk support and target distance on postural adjustments prior to a rapid reaching task by seated subjects. Arch Phys Med Rehabil. 72, 9, 638-641.

Moore, S., Brunt, D., Nesbitt, M. L., Juarez, T. (1992) Investigation of evidence for anticipatory postural adjustments in seated subjects who performed a reaching task. Phys Ther. 72, 5, 335-343.

Mortimer, M., Pernold, G., Wiktorin, C. (2006) Low back pain in a general population. Natural course and influence of physical exercise--a 5-year follow-up of the Musculoskeletal Intervention Center-Norrtaälje Study. Spine (Phila Pa 1976). 31, 26, 3045-3051.

Mount, J. (1996) Effect of practice of a throwing skill in one body position on performance of the skill in an alternate position. Percept Mot Skills. 83, 3 Pt 1, 723-732.

Muraki, S., Akune, T., Oka, H., Mabuchi, A., En-Yo, Y., Yoshida, M., Saika, A., Nakamura, K., Kawaguchi, H., Yoshimura, N. (2009) Association of occupational activity with radiographic knee osteoarthritis and lumbar spondylosis in elderly patients of population-based cohorts: a large-scale population-based study. Arthritis Rheum. 61, 6, 779-786.

Olson, M. W., Li, L., Solomonow, M. (2004) Flexion-relaxation response to cyclic lumbar flexion. Clin Biomech (Bristol, Avon). 19, 8, 769-776.

Olson, M. W., Li, L., Solomonow, M. (2009) Interaction of viscoelastic tissue compliance with lumbar muscles during passive cyclic flexion-extension. J Electromyogr Kinesiol. 19, 1, 30-38.

Olson, M., Solomonow, M., Li, L. (2006) Flexion-relaxation response to gravity. J Biomech. 39, 14, 2545-2554.

Ortiz, D. J., Marcus, M., Gerr, F., Jones, W., Cohen, S. (1997) Measurement variability in upper extremity posture among VDT users. Applied Ergonomics. 28, 2, 139-143.

O'Sullivan, P. B., Burnett, A., Floyd, A. N., Gadsdon, K., Logiudice, J., Miller, D., Quirke, H. (2003) Lumbar repositioning deficit in a specific low back pain population. Spine (Phila Pa 1976). 28, 10, 1074-1079.

O'Sullivan, P. B., Dankaerts, W., Burnett, A. F., Farrell, G. T., Jefford, E., Naylor, C. S., O'Sullivan, K. J. (2006) Effect of different upright sitting postures on spinal-pelvic curvature and trunk muscle activation in a pain-free population. Spine (Phila Pa 1976). 31(19), E707-712.

O'Sullivan, P. B., Mitchell, T., Bulich, P., Waller, R., Holte, J. (2006) The relationship between posture and back muscle endurance in industrial workers with flexion-related low back pain. Man Ther. 11, 4, 264-271.

O'Sullivan, P., Dankaerts, W., Burnett, A., Chen, D., Booth, R., Carlsen, C., Schultz, A. (2006) Evaluation of the flexion relaxation phenomenon of the trunk muscles in sitting. Spine (Phila PA 1976). 31, 17, 2009-2016.

Panjabi, M. M. (1992) The stabilizing system of the spine. Part i. Function, dysfunction, adaptation, and enhancement. J Spinal Disord. 5(4), 383-389, discussion 397.

Panjabi, M., Yamamoto, I., Oxland, T., Crisco, J. (1989) How does posture affect coupling in the lumbar spine? Spine (Phila Pa 1976). 14, 9, 1002-1011.

Parakkat, J., Yang, G., Chany, A. M., Burr, D., Marras, W. S. (2007) The influence of lift frequency, lift duration and work experience on discomfort reporting. *Ergonomics*. 50, 3, 396-409.

Pel, J. J., Spoor, C. W., Pool-Goudzwaard, A. L., Hoek van Dijke, G. A., Snijders, C. J. (2008) Biomechanical analysis of reducing sacroiliac joint shear load by optimization of pelvic muscle and ligament forces. *Ann Biomed Eng*. 36, 3, 415-424.

Petrie, S., Collins, J., Solomonow, M., Wink, C., Chuinard, R. (1997) Mechanoreceptors in the palmar wrist ligaments. *J Bone Joint Surg Br*. 79, 3, 494-496.

Pope, M. H., Goh, K. L., Magnusson, M. L. (2002) Spine ergonomics. *Annu Rev Biomed Eng*. 4, 49-68.

Panjabi, M. and White, A. (2001) Biomechanics in the Musculoskeletal System. Churchill Livingstone.

Reinecke, S. M., Hazard, R. G. (1994a) Continuous passive lumbar motion in seating. Lueder, R. and Noro, K. (Ed): Hard Facts about Soft Machines, The Science of Seating, Philadelphia and London. Taylor and Francis. 141-148.

Richardson, C. A., Snijders, C. J., Hides, J. A., Damen, L., Pas, M. S., Storm, J. (2002) The Relation Between the Transversus Abdominis Muscles, Sacroiliac Joint Mechanics, and Low Back Pain. *Spine*. 27, 4, 399-405.

Rogers, E. L., Granata, K. P. (2006) Disturbed paraspinal reflex following prolonged flexion-relaxation and recovery. *Spine* (Phila Pa 1976). 31, 7, 839-845.

Rohlmann, A., Arntz, U., Graichen, F., Bergmann, G. (2001) Loads on an internal spinal fixation device during sitting. *J Biomechanics*. 34, 8, 989-993.

Rohlmann, A., Bergmann, G., Graichen, F. (1997) Loads on an internal spinal fixation device during walking. *J Biomechanics*. 30, 1, 41-47.

Rohlmann, A., Graichen, F., Bergmann, G. (2002) Loads on an internal spinal fixation device during physical therapy. *Phys Ther*. 82, 1, 44-52.

Rohlmann, A., Zander, T., Schmidt, H., Wilke, H. J., Bergmann, G. (2006) Analysis of the influence of disc degeneration on the mechanical behaviour of a lumbar motion segment using the finite element method. *J Biomech*. 39, 13, 2484-2490.

Rohlmann, A., Claes, L. E., Bergmann, G., Graichen, F., Neef, P., Wilke, H. J. (2001) Comparison of intradiscal pressures and spinal fixator loads for different body positions and exercises. *Ergonomics*. 44, 8, 781-794.

Rohlmann, A., Graichen, F., Weber, U., Bergmann, G. (2000) 2000 Volvo Award winner in biomechanical studies: Monitoring in vivo implant loads with a telemeterized internal spinal fixation device. *Spine* (Phila Pa 1976). 25, 23, 2981-2986.

Saur, Petra M. M., Ensink, Franz-Bernhard M., Frese, Knut, Seeger, Dagmar, Hildebrandt, Jan (1996) Lumbar Range of Motion: Reliability and Validity of the Inclinator Technique in the Clinical Measurement of Trunk Flexibility. Spine. 21, 11, 1332-1338.

Sbriccoli, P., Solomonow, M., Zhou, B. H., Baratta, R. V., Lu, Y., Zhu, M. P., Burger, E. L. (2004) Static load magnitude is a risk factor in the development of cumulative low back disorder. Muscle Nerve. 29, 2, 300-308.

Sbriccoli, P., Yousuf, K., Kupershtein, I., Solomonow, M., Zhou, B. H., Zhu, M. P., Lu, Y. (2004) Static load repetition is a risk factor in the development of lumbar cumulative musculoskeletal disorder. Spine (Phila Pa 1976). 29, 23, 2643-2653.

Scannell, J. P., McGill, S. M. (2003) Lumbar posture--should it, and can it, be modified? A study of passive tissue stiffness and lumbar position during activities of daily living. Phys Ther. 83, 10, 907-917.

Seidler, A, Bolm-Audorff, U, Heiskel, H, Henkel, N, Roth-Küver, B, Kaiser, U, Bickeböller, R, Willingstorfer, W J, Beck, W, Elsner, G (2001) The role of cumulative physical work load in lumbar spine disease: risk factors for lumbar osteochondrosis and spondylosis associated with chronic complaints. Occupational and Environmental Medicine. 58, 11, 735-746.

Seidler, A., Bolm-Audorff, U., Siol, T., Henkel, N., Fuchs, C., Schug, H., Leheta, F., Marquardt, G., Schmitt, E., Ulrich, P. T., Beck, W., Missalla, A., Elsner, G. (2003) Occupational risk factors for symptomatic lumbar disc herniation, a case-control study. Occup Environ Med. 60, 11, 821-830.

Smith, A., O'Sullivan, P., Straker, L. (2008) Classification of sagittal thoraco-lumbo-pelvic alignment of the adolescent spine in standing and its relationship to low back pain. Spine (Phila Pa 1976). 33, 19, 2101-2107.

Snijders, C. J., Goossens, R. H. M., van Dijke, G. A. Hoek (2000) Minimization of Pressure and Shear Load in Sitting and Lying, Based on Biomechanical Modeling. Human Factors and Ergonomics Society Annual Meeting Proceedings. 44, 692-695.

Snijders, C. J., Hermans, P. F., Niesing, R., Jan Kleinrensink, G., Pool-Goudzwaard, A. (2008) Effects of slouching and muscle contraction on the strain of the iliolumbar ligament. Man Ther. 13, 4, 325-333.

Snijders, C. J., Hermans, P. F., Niesing, R., Spoor, C. W., Stoeckart, R. (2004) The influence of slouching and lumbar support on iliolumbar ligaments, intervertebral discs and sacroiliac joints. Clin Biomech (Bristol, Avon). 19, 4, 323-329.

Soderberg, G. L., Blanco, M. K., Cosentino, T. L., Kurdelmeier, K. A. (1986) An EMG analysis of posterior trunk musculature during flat and anteriorly inclined sitting. Hum Factors. 28, 4, 483-491.

Solomonow, D., Davidson, B., Zhou, B. H., Lu, Y., Patel, V., Solomonow, M. (2008) Neuromuscular neutral zones response to cyclic lumbar flexion. J Biomech. 41, 13, 2821-2828.

Solomonow, M. (2001) Knee bracing after ACL reconstruction. Arch Phys Med Rehabil. 82, 5, 709-710.

Solomonow, M. (2004) Ligaments: a source of work-related musculoskeletal disorders. J Electromyogr Kinesiol. 14, 1, 49-60.

Solomonow, M. (2006) Sensory-motor control of ligaments and associated neuromuscular disorders. J Electromyogr Kinesiol. 16, 6, 549-567.

Solomonow, M., Baratta, R. V., Zhou, B. H., Burger, E., Zieske, A., Gedalia, A. (2003b) Muscular dysfunction elicited by creep of lumbar viscoelastic tissue. J Electromyogr Kinesiol. 13, 4, 381-396.

Solomonow, M., Eversull, E., He Zhou, B., Baratta, R. V., Zhu, M. P. (2001) Neuromuscular neutral zones associated with viscoelastic hysteresis during cyclic lumbar flexion. Spine (Phila Pa 1976). 26, 14, E314-324.

Solomonow, M., Krogsgaard, M. (2001) Sensorimotor control of knee stability. A review. Scand J Med Sci Sports. 11, 2, 64-80.

Solomonow, M., Lewis, J. (2002) Reflex from the ankle ligaments of the feline. J Electromyogr Kinesiol. 12, 3, 193-198.

Solomonow, M., Zhou, B. H., Baratta, R. V., Burger, E. (2003) Biomechanics and electromyography of a cumulative lumbar disorder: response to static flexion. Clin Biomech (Bristol, Avon). 18, 10, 890-898.

Solomonow, M., Zhou, B., Baratta, R. V., Zhu, M., Lu, Y. (2002) Neuromuscular disorders associated with static lumbar flexion: a feline model. J Electromyogr Kinesiol. 12, 2, 81-90.

Szeto, G.P., Straker, L., Raine, S. (2002) A field comparison of neck and shoulder postures in symptomatic and asymptomatic office workers. Applied Ergonomics. 33(1), 75-84.

Takayanagi, K., Takahashi, K., Yamagata, M., Moriya, H., Kitahara, H., Tamaki, T. (2001) Using Cineradiography for Continuous Dynamic-Motion Analysis of the Lumbar Spine. Spine. 26, 17, 1858-1865.

Thompson, Rosemary Elizabeth (2002) Mechanical Effects of Degeneration in Lumbar Intervertebral Discs. PhD Doctorate. Queensland University of Technology. Faculty of Built Environment and Engineering.

Tichauer, E.R. (1978) The Biochemical Basis of Ergonomics: Anatomy Applied to the Design of Work Situation. Wiley, John & Sons.

Tomasek, J. J., Haaksma, C. J., Schwartz, R. J., Vuong, D. T., Zhang, S. X., Ash, J. D., Ma, J. X., Al-Ubaidi, M. R. (2006) Deletion of smooth muscle alpha-actin alters blood-retina barrier permeability and retinal function. Invest Ophthalmol Vis Sci. 47, 6, 2693-2700.

Tomasek, J. J., Vaughan, M. B., Haaksma, C. J. (1999) Cellular structure and biology of Dupuytren's disease. Hand Clin. 15, 1, 21-34.

Van Deursen, D.L., Snijders, C.J., Van Dieen, J.H., Kingma, I., Van Deursen, L.L. (2001) The effect of passive vertebral rotation on pressure in the nucleus pulposus. J Biomech. 34(3), 405-408.

Van Deursen, D.L., Lengsfeld, M., Snijders, C.J., Evers, J.J., Goossens, R.H. (2000) Mechanical effects of continuous passive motion on the lumbar spine in seating. J Biomech. 33(6), 695-699.

van Deursen, D. L., Snijders, C. J., van Deursen, L. L. J. M. (2000) Sitting with Rotary Continuous Passive Motion Biomechanical explanation of health effects. Human Factors and Ergonomics Society Annual Meeting Proceedings. 44, 272-275.

van Deursen, L L. J. M., van Deursen, D. L., Patijn, J. (2000) Sitting and Low Back Pain Sitting with Rotary Continuous Passive Motion A New Seating Concept. Human Factors and Ergonomics Society Annual Meeting Proceedings. 44, 406-406.

van Dieen, J. H., Jansen, S. M., Housheer, A. F. (1997) Differences in low back load between kneeling and seated working at ground level. Applied Ergonomics. 28, 5-6, 355-363.

van Dieen, J. H., Jansen, S. M., Housheer, A. F. (1997) Differences in low back load between kneeling and seated working at ground level. Applied Ergonomics. 28, 5-6, 355-363.

van Veelen, Martine A., Goossens, Richard H. M., Chang, Annemarie, Kumar, Shrawan, Snijders, Chris J. (2000) High Risks of High Backrests. Human Factors and Ergonomics Society Annual Meeting Proceedings. 44, 696-699.

Vergara, M., Page, A. (2000) System to measure the use of the backrest in sitting-posture office tasks. Applied Ergonomics. 31, 3, 247-254.

Vergara, M., Page, A. (2000) Technique to measure lumbar curvature in the ergonomic evaluation of chairs: description and validation. Clin Biomech (Bristol, Avon). 15, 10, 786-789.

Vergara, M., Page, A. (2002) Relationship between comfort and back posture and mobility in sitting-posture. Applied Ergonomics. 33, 1, 1-8.

Vingard, E., Mortimer, M., Wiktorin, C., Pernold, R. P. T. G., Fredriksson, K., Nemeth, G., Alfredsson, L. (2002) Seeking care for low back pain in the general population: a two-year follow-up study: results from the MUSIC-Norrtälje Study. Spine (Phila Pa 1976). 27, 19, 2159-2165.

Walsh, E. G. (2003) Axial rotation of the lower human spine by rhythmic torques automatically generated at the resonant frequency. Exp Physiol. 88, 3, 305-308.

Whistance, R. S., Adams, L. P., van Geems, B. A., Bridger, R. S. (1995) Postural adaptations to workbench modifications in standing workers. Ergonomics. 38, 12, 2485-2503.

Wilder, D., Magnusson, M. L., Fenwick, J., Pope, M. (1994) The effect of posture and seat suspension design on discomfort and back muscle fatigue during simulated truck driving. Applied Ergonomics. 25, 2, 66-76.

Wilke, H. J., Rohlmann, A., Neller, S., Graichen, F., Claes, L., Bergmann, G. (2003) ISSLS prize winner: A novel approach to determine trunk muscle forces during flexion and extension: a comparison of data from an in vitro experiment and in vivo measurements. Spine. 28, 23, 2585-2593.

Wilke, H., Neef, P., Hinz, B., Seidel, H., Claes, L. (2001) Intradiscal pressure together with anthropometric data--a data set for the validation of models. Clin Biomech (Bristol, Avon). 16 Suppl 1, S111-126.

Williams, G. N., Higgins, M. J., Lewek, M. D. (2002) Aging skeletal muscle: physiologic changes and the effects of training. Phys Ther. 82, 1, 62-68.

Williams, M., Solomonow, M., Zhou, B. H., Baratta, R. V., Harris, M. (2000) Multifidus spasms elicited by prolonged lumbar flexion. Spine (Phila Pa 1976). 25, 22, 2916-2924.

Wolinsky, F. D., Miller, D. K., Andresen, E. M., Malmstrom, T. K., Miller, J. P. (2005) Further evidence for the importance of subclinical functional limitation and subclinical disability assessment in gerontology and geriatrics. J Gerontol B Psychol Sci Soc Sci. 60, 3, S146-151.

Womersley, L., May, S. (2006) Sitting posture of subjects with postural backache. J Manipulative Physiol Ther. 29, 3, 213-218.

Yang, G., Chany, A. M., Parakkat, J., Burr, D., Marras, W. S. (2007) The effects of work experience, lift frequency and exposure duration on low back muscle oxygenation. Clin Biomech (Bristol, Avon). 22, 1, 21-27.

Youdas, J. W., Garrett, T. R., Egan, K. S., Therneau, T. M. (2000) Lumbar lordosis and pelvic inclination in adults with chronic low back pain. Phys Ther. 80, 3, 261-275.

Youssef, J., Davidson, B., Zhou, B. H., Lu, Y., Patel, V., Solomonow, M. (2008) Neuromuscular neutral zones response to static lumbar flexion: muscular stability compensator. Clin Biomech (Bristol, Avon). 23, 7, 870-880.

Zhou, B. H., Baratta, R. V., Solomonow, M., Zhu, M., Lu, Y. (2000) Closed-loop control of muscle length through motor unit recruitment in load-moving conditions. J Biomech. 33, 7, 827-835.

About Rani Lueder, CPE

Rani Lueder, MSIE, CPE is President of Humanics ErgoSystems, Inc. an ergonomics consulting firm in Encino, California. She has consulted in workplace ergonomics, product design research and accommodating special populations since 1982.

She consulted for corporations, governments and universities in nine countries. Rani was a member of the American National Standards Institute committee responsible for revising ANSI guidelines for seating and computer workstation design and is currently participating as part of the US team of experts on the International Organization for Standardization guideline ISO/TC 159/WG 2, "Ergonomics for people with special requirements" in standards development.

Since 1988, she continues to serve on retainer to several organizations in Japan, including the Waseda University's Seating Research Lab, ErgoSeating Japan and the Japan Institute of Human Posture Research. She served as US organizing chair for the Second International Conference on Seated Posture, held in Tokyo.

She recently co-edited her third book on ergonomics, this one on ergonomics for children. Her second edited / co-authored book, "Hard Facts about Soft Machines: The ergonomics of seating" (Dec. 1995) is available internationally from Taylor and Francis (London). Previously, she edited and co-authored the book "The Ergonomics Payoff, Designing the electronic office" (Holt Reinhart and Winston).

She has an MSIE in Ergonomics/Industrial Engineering from Virginia Tech and is a member of HFES (US). The Board of Certification certifies her in Professional Ergonomics. She serves on the Advisory Board of two national ergonomics conferences.

More details about her activities are at www.humanics-es.com